

ABIRATERONE

Products Affected

• abiraterone acetate

PA Criteria	Criteria Details
Exclusion Criteria	None
Required Medical Information	Diagnosis of one of the following A.) Castration-resistant metastatic prostate cancer (CRPC), or B.) High risk, castration-sensitive metastatic prostate cancer (CSPC). For treatment of CRPC and CSPC, abiraterone will be used in combination with prednisone AND one of the following applies 1.) Used in combination with a gonadotropin-releasing hormone (GnRH) analog (e.g. leuprolide, triptorelin), OR 2) Patient has received bilateral orchiectomy
Age Restrictions	18 years of age and older
Prescriber Restrictions	Prescribed by or in consultation with an oncologist or urologist
Coverage Duration	12 months
Other Criteria	None
Indications	All Medically-accepted Indications.
Off-Label Uses	N/A
Part B Prerequisite	No

Formulary ID: 24449 version 13

Last Updated: 06/20/2024 Effective: 07/01/2024



ACITRETIN

Products Affected

• acitretin

PA Criteria	Criteria Details
Exclusion Criteria	Any of the following A.) Severely impaired liver or kidney function, B.) Chronic abnormally elevated blood lipid values, C.) Concomitant use of methotrexate or tetracyclines, D.) Pregnancy
Required Medical Information	Diagnosis of severe, recalcitrant psoriasis (including plaque, guttate, erythrodermic palmar- plantar and pustular) AND patient must have tried and failed, contraindication or intolerance to one formulary first line agent (e.g., Topical Corticosteroids (betamethasone, fluocinonide, desoximetasone), Topical Calcipotriene/Calcitriol, Topical Calcipotriene, OR Topical Tazarotene)
Age Restrictions	18 years of age and older
Prescriber Restrictions	Prescribed by or in consultation with a dermatologist
Coverage Duration	12 months
Other Criteria	None
Indications	All Medically-accepted Indications.
Off-Label Uses	N/A
Part B Prerequisite	No

Formulary ID: 24449 version 13

Last Updated: 06/20/2024 Effective: 07/01/2024



ACTIMMUNE

Products Affected

• ACTIMMUNE

PA Criteria	Criteria Details
Exclusion Criteria	None
Required Medical Information	Diagnosis of one of the following A.) Chronic granulomatous disease for use in reducing the frequency and severity of serious infections, or B.) Severe, malignant osteopetrosis (SMO)
Age Restrictions	None
Prescriber Restrictions	None
Coverage Duration	12 months
Other Criteria	B vs D determination required per CMS guidance
Indications	All Medically-accepted Indications.
Off-Label Uses	N/A
Part B Prerequisite	No

Formulary ID: 24449 version 13

Last Updated: 06/20/2024 Effective: 07/01/2024



ADEMPAS

Products Affected

ADEMPAS

PA Criteria	Criteria Details
Exclusion Criteria	Any of the following A.) Concomitant administration with nitrates or nitric oxide donors (such as amyl nitrate) in any form, B.) Concomitant administration with phosphodiesterase inhibitors, including specific PDE-5 inhibitors (such as sildenafil, tadalafil, or vardenafil) or non-specific PDE inhibitors (such as dipyridamole or theophylline), C.) Pregnancy, or D.) Patients with pulmonary hypertension associated with idiopathic interstitial pneumonia
Required Medical Information	Diagnosis of one of the following A.) Pulmonary arterial hypertension (WHO group I) and diagnosis was confirmed by right heart catheterization or Doppler echocardiogram if patient is unable to undergo a right heart catheterization (e.g., patient is frail, elderly, etc.), or B.) Chronic thromboembolic pulmonary hypertension (CTEPH, WHO group 4) and patient has persistent or recurrent disease after surgical treatment (e.g., pulmonary endarterectomy) or has CTEPH that is inoperable (Female patients must be enrolled in the ADEMPAS REMS program)
Age Restrictions	18 years of age and older
Prescriber Restrictions	Prescribed by or in consultation with a cardiologist or pulmonologist
Coverage Duration	Initial: 6 months, Renewal: 12 months
Other Criteria	None
Indications	All Medically-accepted Indications.
Off-Label Uses	N/A
Part B Prerequisite	No

Formulary ID: 24449 version 13 Last Updated: 06/20/2024

Effective: 07/01/2024



AKEEGA

Products Affected

AKEEGA

PA Criteria	Criteria Details
Exclusion Criteria	None
Required Medical Information	Diagnosis of deleterious or suspected deleterious BRCA-mutated (BRCAm) metastatic castration-resistant prostate cancer (mCRPC) AND used in combination with prednisone
Age Restrictions	18 years or older
Prescriber Restrictions	Prescribed by or in consultation with an oncologist
Coverage Duration	12 months
Other Criteria	None
Indications	All Medically-accepted Indications.
Off-Label Uses	N/A
Part B Prerequisite	No

Formulary ID: 24449 version 13 Last Updated: 06/20/2024

Effective: 07/01/2024



ALECENSA

Products Affected

• ALECENSA

PA Criteria	Criteria Details
Exclusion Criteria	None
Required Medical Information	Diagnosis of metastatic anaplastic lymphoma kinase (ALK) positive non-small cell lung cancer as detected by an FDA-approved test
Age Restrictions	18 years of age and older
Prescriber Restrictions	None
Coverage Duration	12 months
Other Criteria	None
Indications	All Medically-accepted Indications.
Off-Label Uses	N/A
Part B Prerequisite	No

Formulary ID: 24449 version 13

Last Updated: 06/20/2024 Effective: 07/01/2024



ALPHA-1 PROTEINASE INHIBITOR

Products Affected

• PROLASTIN-C INTRAVENOUS SOLUTION

PA Criteria	Criteria Details
Exclusion Criteria	Immunoglobulin A (IgA) deficiency with antibodies against IgA
Required Medical Information	Diagnosis of alpha-1 proteinase inhibitor (alpha-1 antitrypsin) deficiency in adult patients with emphysema
Age Restrictions	18 years of age and older
Prescriber Restrictions	Prescribed by or in consultation with a pulmonologist
Coverage Duration	12 months
Other Criteria	B vs D determination required per CMS guidance
Indications	All Medically-accepted Indications.
Off-Label Uses	N/A
Part B Prerequisite	No

Formulary ID: 24449 version 13 Last Updated: 06/20/2024

Effective: 07/01/2024



ALUNBRIG

Products Affected

• ALUNBRIG

PA Criteria	Criteria Details
Exclusion Criteria	None
Required Medical Information	Diagnosis of anaplastic lymphoma kinase-positive (ALK) metastatic non-small cell lung cancer (NSCLC)
Age Restrictions	18 years of age and older
Prescriber Restrictions	Prescribed by or in consultation with an oncologist or hematologist
Coverage Duration	12 months
Other Criteria	None
Indications	All Medically-accepted Indications.
Off-Label Uses	N/A
Part B Prerequisite	No

Formulary ID: 24449 version 13

Last Updated: 06/20/2024 Effective: 07/01/2024



ALVAIZ

Products Affected

ALVAIZ

PA Criteria	Criteria Details
Exclusion Criteria	None
Required Medical Information	Diagnosis of one of the following A.) Chronic idiopathic thrombocytopenic purpura (ITP), B.) Chronic hepatitis C infection associated thrombocytopenia, or C.) Severe aplastic anemia with insufficient response to immunosuppressive therapy
Age Restrictions	None
Prescriber Restrictions	None
Coverage Duration	12 months
Other Criteria	None
Indications	All Medically-accepted Indications.
Off-Label Uses	N/A
Part B Prerequisite	No

Formulary ID: 24449 version 13

Last Updated: 06/20/2024 Effective: 07/01/2024



AMBRISENTAN

Products Affected

• ambrisentan

PA Criteria	Criteria Details
1 A CHICHA	Criteria Details
Exclusion Criteria	Any of the following A.) Pregnancy, or B.) Idiopathic pulmonary fibrosis (IPF), including those with pulmonary hypertension
Required Medical Information	Diagnosis of pulmonary arterial hypertension classified as WHO Group I, confirmed by right heart catheterization or Doppler echocardiogram if patient is unable to undergo a right heart catheterization (e.g., patient is frail, elderly, etc.)
Age Restrictions	18 years of age and older
Prescriber Restrictions	Prescribed by or in consultation with a cardiologist or pulmonologist
Coverage Duration	Initial: 6 months, Renewal: 12 months
Other Criteria	None
Indications	All Medically-accepted Indications.
Off-Label Uses	N/A
Part B Prerequisite	No

Formulary ID: 24449 version 13

Last Updated: 06/20/2024 Effective: 07/01/2024



ARCALYST

Products Affected

ARCALYST

PA Criteria	Criteria Details
Exclusion Criteria	None
Required Medical Information	Diagnosis of one of the following A.) Cryopyrin-associated periodic syndromes (CAPS), including familial cold autoinflammatory syndrome (FCAS) and Muckle-Wells Syndrome (MWS), B.) Deficiency of interleukin-1 receptor antagonist (DIRA) and patient requires maintenance therapy for remission, or C.) Recurrent pericarditis (RP) and reduction in risk of recurrence
Age Restrictions	None
Prescriber Restrictions	None
Coverage Duration	12 months
Other Criteria	B vs D determination required per CMS guidance
Indications	All Medically-accepted Indications.
Off-Label Uses	N/A
Part B Prerequisite	No

Formulary ID: 24449 version 13

Last Updated: 06/20/2024 Effective: 07/01/2024



ARIKAYCE

Products Affected

ARIKAYCE

PA Criteria	Criteria Details
Exclusion Criteria	Known sensitivity to any aminoglycoside
Required Medical Information	Diagnosis of pulmonary Mycobacterium avium complex (MAC) infection and used as part of a combination antibacterial regimen in treatment refractory patients (greater than 6 months of a multidrug background regimen)
Age Restrictions	18 years of age and older
Prescriber Restrictions	Prescribed by or in consultation with an infectious disease specialist or pulmonologist
Coverage Duration	12 months
Other Criteria	None
Indications	All Medically-accepted Indications.
Off-Label Uses	N/A
Part B Prerequisite	No

Formulary ID: 24449 version 13

Last Updated: 06/20/2024 Effective: 07/01/2024



AUGTYRO

Products Affected

AUGTYRO

PA Criteria	Criteria Details
Exclusion Criteria	None
Required Medical Information	Diagnosis of locally advanced or metastatic ROS1-positive non-small cell lung cancer
Age Restrictions	18 years or older
Prescriber Restrictions	Prescribed by or in consultation with an oncologist
Coverage Duration	12 months
Other Criteria	None
Indications	All Medically-accepted Indications.
Off-Label Uses	N/A
Part B Prerequisite	No

Formulary ID: 24449 version 13 Last Updated: 06/20/2024

Effective: 07/01/2024



AURYXIA

Products Affected

AURYXIA

PA Criteria	Criteria Details
Exclusion Criteria	Iron overload syndrome (e.g. hemochromatosis)
Required Medical Information	Diagnosis of hyperphosphatemia in patients with chronic kidney disease (CKD) on dialysis
Age Restrictions	18 years of age and older
Prescriber Restrictions	Prescribed by or in consultation with a hematologist or nephrologist
Coverage Duration	12 months
Other Criteria	Ferric Citrate is NOT approvable for iron deficiency anemia per Part D law
Indications	All Medically-accepted Indications.
Off-Label Uses	N/A
Part B Prerequisite	No

Formulary ID: 24449 version 13 Last Updated: 06/20/2024

Effective: 07/01/2024



AUSTEDO

Products Affected

- AUSTEDO
- AUSTEDO XR

• AUSTEDO XR PATIENT TITRATION

PA Criteria	Criteria Details
Exclusion Criteria	Any of the following A.) Suicidal ideation and/or untreated or inadequately treated depression in a patient with Huntington's Disease, B.) Hepatic impairment, C.) Concomitant use of MAOIs, reserpine, tetrabenazine, or valbenazine
Required Medical Information	Diagnosis of one of the following A.) Chorea associated with Huntington's disease (Huntington's chorea), or B.) Tardive dyskinesia
Age Restrictions	18 years of age and older
Prescriber Restrictions	Prescribed by or in consultation with a neurologist or psychiatrist
Coverage Duration	12 months
Other Criteria	None
Indications	All Medically-accepted Indications.
Off-Label Uses	N/A
Part B Prerequisite	No

Formulary ID: 24449 version 13 Last Updated: 06/20/2024

Effective: 07/01/2024



AYVAKIT

Products Affected

AYVAKIT

PA Criteria	Criteria Details
Exclusion Criteria	None
Required Medical Information	Diagnosis of one of the following A.) Unresectable or metastatic gastrointestinal stromal tumor, with a platelet-derived growth factor receptor alpha (PDGFRA) exon 18 mutation, including PDGFRA D842V mutations, B.) Advanced Systemic Mastocytosis (AdvSM), including aggressive systemic mastocytosis (ASM), systemic mastocytosis with an associated hematological neoplasm (SMAHN), or mast cell leukemia (MCL), and platelet count of at least 50,000/mcL, or C.) Indolent systemic mastocytosis
Age Restrictions	18 years of age and older
Prescriber Restrictions	None
Coverage Duration	12 months
Other Criteria	None
Indications	All Medically-accepted Indications.
Off-Label Uses	N/A
Part B Prerequisite	No

Formulary ID: 24449 version 13

Last Updated: 06/20/2024 Effective: 07/01/2024



BALVERSA

Products Affected

• BALVERSA

PA Criteria	Criteria Details
Exclusion Criteria	None
Required Medical Information	Diagnosis of locally advanced or metastatic urothelial carcinoma and both of the following 1.) Susceptible fibroblast growth factor receptor (FGFR)3 or FGFR2 genetic alterations confirmed by an FDA-approved diagnostic test, and 2.) Patient has progressed during or following at least one line of prior platinum-containing chemotherapy, including within 12 months of neoadjuvant or adjuvant platinum-containing chemotherapy
Age Restrictions	18 years of age and older
Prescriber Restrictions	Prescribed by or in consultation with an oncologist or urologist
Coverage Duration	12 months
Other Criteria	None
Indications	All Medically-accepted Indications.
Off-Label Uses	N/A
Part B Prerequisite	No

Formulary ID: 24449 version 13

Last Updated: 06/20/2024 Effective: 07/01/2024



BENLYSTA

Products Affected

• BENLYSTA SUBCUTANEOUS

PA Criteria	Criteria Details
Exclusion Criteria	None
Required Medical Information	Diagnosis of one of the following A.) Active, autoantibody-positive, system lupus erythematosus (SLE), or B.) Active lupus nephritis and patient is receiving standard therapy
Age Restrictions	None
Prescriber Restrictions	Prescribed by or in consultation with a nephrologist or rheumatologist
Coverage Duration	12 months
Other Criteria	None
Indications	All Medically-accepted Indications.
Off-Label Uses	N/A
Part B Prerequisite	No

Formulary ID: 24449 version 13

Last Updated: 06/20/2024 Effective: 07/01/2024



BESREMI

Products Affected

• BESREMI

PA Criteria	Criteria Details
Exclusion Criteria	Any of the following A.) Existence of, or history of severe psychiatric disorders (severe depression, suicidal ideation, or suicide attempt), B.) Hypersensitivity to interferons including interferon alfa-2b or excipients, C.) Hepatic impairment (Child-Pugh B or C), D.) History or presence of active serious or untreated autoimmune disease, or E.) Immunosuppressed transplant recipients
Required Medical Information	Diagnosis of polycythemia vera
Age Restrictions	18 years of age and older
Prescriber Restrictions	Prescribed by or in consultation with an oncologist or hematologist
Coverage Duration	12 months
Other Criteria	None
Indications	All Medically-accepted Indications.
Off-Label Uses	N/A
Part B Prerequisite	No

Formulary ID: 24449 version 13 Last Updated: 06/20/2024

Effective: 07/01/2024



BEXAROTENE GEL

Products Affected

• bexarotene external

PA Criteria	Criteria Details
Exclusion Criteria	None
Required Medical Information	Diagnosis of primary cutaneous T-cell lymphoma (CTCL Stage 1A/1B) and patient had an inadequate response, is intolerant to, or has a contraindication to at least one prior systemic therapy (e.g., corticosteroids) indicated for cutaneous manifestations of CTCL
Age Restrictions	None
Prescriber Restrictions	Prescribed by or in consultation with an oncologist or dermatologist
Coverage Duration	12 months
Other Criteria	None
Indications	All Medically-accepted Indications.
Off-Label Uses	N/A
Part B Prerequisite	No

Formulary ID: 24449 version 13 Last Updated: 06/20/2024

Effective: 07/01/2024



BEXAROTENE ORAL

Products Affected

• bexarotene oral

PA Criteria	Criteria Details
Exclusion Criteria	Pregnancy
Required Medical Information	Diagnosis of cutaneous T-cell lymphoma (CTCL) and patient is not a candidate for or had an inadequate response, is intolerant to, or has a contraindication to at least one prior systemic therapy (e.g., corticosteroids) for cutaneous manifestations of CTCL
Age Restrictions	None
Prescriber Restrictions	Prescribed by or in consultation with an dermatologist, hematologist, or oncologist
Coverage Duration	12 months
Other Criteria	None
Indications	All Medically-accepted Indications.
Off-Label Uses	N/A
Part B Prerequisite	No

Formulary ID: 24449 version 13 Last Updated: 06/20/2024

Effective: 07/01/2024



BOSENTAN

Products Affected

• bosentan

PA Criteria	Criteria Details
Exclusion Criteria	Any of the following A.) Concomitant cyclosporine A or glyburide therapy, or B.) Pregnancy
Required Medical Information	Diagnosis of pulmonary arterial hypertension (WHO Group I) and patient has New York Heart Association (NYHA) Functional Class II-IV, confirmed by right heart catheterization or Doppler echocardiogram if patient is unable to undergo a right heart catheterization (e. g., patient is frail, elderly, etc.)
Age Restrictions	None
Prescriber Restrictions	Prescribed by or in consultation with a cardiologist or pulmonologist
Coverage Duration	Initial: 6 months, Renewal: 12 months
Other Criteria	None
Indications	All Medically-accepted Indications.
Off-Label Uses	N/A
Part B Prerequisite	No

Formulary ID: 24449 version 13 Last Updated: 06/20/2024

Effective: 07/01/2024



BOSULIF

Products Affected

• BOSULIF

PA Criteria	Criteria Details
Exclusion Criteria	None
Required Medical Information	Diagnosis of one of the following A.) Chronic, accelerated, or blast phase Philadelphia chromosome-positive (Ph+) chronic myelogenous leukemia (CML) with resistance or inadequate response to prior therapy, or B.) Newly diagnosed chronic phase Philadelphia chromosome-positive (Ph+) CML
Age Restrictions	None
Prescriber Restrictions	Prescribed by or in consultation with an oncologist or hematologist
Coverage Duration	12 months
Other Criteria	None
Indications	All Medically-accepted Indications.
Off-Label Uses	N/A
Part B Prerequisite	No

Formulary ID: 24449 version 13

Last Updated: 06/20/2024 Effective: 07/01/2024



BRAFTOVI

Products Affected

• BRAFTOVI ORAL CAPSULE 75 MG

PA Criteria	Criteria Details
Exclusion Criteria	None
Required Medical Information	Diagnosis of one of the following A.) unresectable or metastatic melanoma with documented BRAF V600E or V600K mutation as detected by a FDA-approved test and used in combination with binimetinib, B.) metastatic colorectal cancer with documented BRAF V600E mutation as detected by a FDA-approved test, patient has received prior therapy, and braftovi used in combination with cetuximab, or C.) Metastatic non-small cell lung cancer with a BRAF V600E mutation as detected by an FDA-approved test and used in combination with binimetinib
Age Restrictions	18 years of age and older
Prescriber Restrictions	Prescribed by or in consultation with an oncologist
Coverage Duration	12 months
Other Criteria	None
Indications	All Medically-accepted Indications.
Off-Label Uses	N/A
Part B Prerequisite	No

Formulary ID: 24449 version 13 Last Updated: 06/20/2024

Effective: 07/01/2024



BRONCHITOL

Products Affected

• BRONCHITOL

PA Criteria	Criteria Details
Exclusion Criteria	None
Required Medical Information	Diagnosis of Cystic fibrosis of the lung
Age Restrictions	None
Prescriber Restrictions	Prescribed by or in consultation with a pulmonologist or prescribing practitioner is from a CF center accredited by the Cystic Fibrosis Foundation
Coverage Duration	12 months
Other Criteria	None
Indications	All Medically-accepted Indications.
Off-Label Uses	N/A
Part B Prerequisite	No

Formulary ID: 24449 version 13

Last Updated: 06/20/2024 Effective: 07/01/2024



BRUKINSA

Products Affected

• BRUKINSA

PA Criteria	Criteria Details
Exclusion Criteria	None
Required Medical Information	Diagnosis of one of the following: A.) mantle cell lymphoma (MCL) and patient has received at least one prior therapy, B.) Treatment of adult patients with Waldenstrom macroglobulinemia, C.) Treatment of adult patients with relapsed or refractory marginal zone lymphoma who have received at least one anti-CD20-based regimen, D.) Chronic lymphocytic leukemia, E.) Small lymphocytic lymphoma, or F.) Relapsed or refractory follicular lymphoma, in combination with obinutuzumab, after 2 or more lines of systemic therapy
Age Restrictions	18 years of age and older
Prescriber Restrictions	None
Coverage Duration	12 months
Other Criteria	None
Indications	All Medically-accepted Indications.
Off-Label Uses	N/A
Part B Prerequisite	No

Formulary ID: 24449 version 13 Last Updated: 06/20/2024

Effective: 07/01/2024



BYLVAY

Products Affected

• BYLVAY

• BYLVAY (PELLETS)

PA Criteria	Criteria Details
Exclusion Criteria	None
Required Medical Information	Diagnosis of one of the following A.) Progressive familial intrahepatic cholestasis-associated pruritus, or B.) Cholestatic pruritus in patients with Alagille syndrome
Age Restrictions	None
Prescriber Restrictions	None
Coverage Duration	12 months
Other Criteria	None
Indications	All Medically-accepted Indications.
Off-Label Uses	N/A
Part B Prerequisite	No

Formulary ID: 24449 version 13

Last Updated: 06/20/2024 Effective: 07/01/2024



CABLIVI

Products Affected

CABLIVI

PA Criteria	Criteria Details
Exclusion Criteria	None
Required Medical Information	Diagnosis of acquired thrombotic thrombocytopenic purpura (aTTP) and used in combination with plasma exchange and immunosuppression therapy
Age Restrictions	18 years of age and older
Prescriber Restrictions	Prescribed by or in consultation with a hematologist or oncologist
Coverage Duration	12 months
Other Criteria	None
Indications	All Medically-accepted Indications.
Off-Label Uses	N/A
Part B Prerequisite	No

Formulary ID: 24449 version 13

Last Updated: 06/20/2024 Effective: 07/01/2024



CABOMETYX

Products Affected

CABOMETYX

PA Criteria	Criteria Details
Exclusion Criteria	None
Required Medical Information	Diagnosis of one of the following A.) Advanced renal cell carcinoma, B.) Advanced hepatocellular carcinoma (HCC) and patient has been previously treated with sorafenib, C.) Advanced renal cell carcinoma and used as first line treatment in combination with nivolumab or D.) treatment of adults and pediatric patients 12 years and older with locally advanced or metastatic differentiated thyroid cancer that has progressed following VEGFR-targeted therapy and who are radioactive iodine-refractory or ineligible
Age Restrictions	None
Prescriber Restrictions	None
Coverage Duration	12 months
Other Criteria	None
Indications	All Medically-accepted Indications.
Off-Label Uses	N/A
Part B Prerequisite	No

Formulary ID: 24449 version 13

Last Updated: 06/20/2024 Effective: 07/01/2024



CALQUENCE

Products Affected

• CALQUENCE

PA Criteria	Criteria Details
Exclusion Criteria	None
Required Medical Information	Diagnosis of one of the following A.) Mantle cell lymphoma (MCL) and patient has received at least 1 prior therapy, B.) Chronic lymphocytic leukemia (CLL), or C.) Small lymphocytic lymphoma (SLL)
Age Restrictions	18 years of age and older
Prescriber Restrictions	Prescribed by or in consultation with an oncologist or hematologist
Coverage Duration	12 months
Other Criteria	None
Indications	All Medically-accepted Indications.
Off-Label Uses	N/A
Part B Prerequisite	No

Formulary ID: 24449 version 13 Last Updated: 06/20/2024

Effective: 07/01/2024



CAMZYOS

Products Affected

CAMZYOS

PA Criteria	Criteria Details
Exclusion Criteria	None
Required Medical Information	Diagnosis of symptomatic New York Heart Association (NYHA) class II-III obstructive hypertrophic cardiomyopathy (HCM) in adult patients
Age Restrictions	18 years of age and older
Prescriber Restrictions	None
Coverage Duration	12 months
Other Criteria	None
Indications	All Medically-accepted Indications.
Off-Label Uses	N/A
Part B Prerequisite	No

Formulary ID: 24449 version 13

Last Updated: 06/20/2024 Effective: 07/01/2024



CAPRELSA

Products Affected

CAPRELSA

PA Criteria	Criteria Details
Exclusion Criteria	Congenital long QT syndrome
Required Medical Information	Diagnosis of metastatic or unresectable locally advanced medullary thyroid cancer (MTC) AND disease is symptomatic or progressive
Age Restrictions	18 years of age and older
Prescriber Restrictions	None
Coverage Duration	12 months
Other Criteria	None
Indications	All Medically-accepted Indications.
Off-Label Uses	N/A
Part B Prerequisite	No

Formulary ID: 24449 version 13

Last Updated: 06/20/2024 Effective: 07/01/2024



CARGLUMIC ACID

Products Affected

• carglumic acid oral tablet soluble

PA Criteria	Criteria Details
Exclusion Criteria	None
Required Medical Information	Diagnosis of one of the following A.) N-acetyl glutamate synthase (NAGS) deficiency (confirmed by appropriate genetic testing) with acute or chronic hyperammonemia, or B.) Propionic or methylmalonic acidemia with acute hyperammonemia
Age Restrictions	None
Prescriber Restrictions	None
Coverage Duration	12 months
Other Criteria	None
Indications	All Medically-accepted Indications.
Off-Label Uses	N/A
Part B Prerequisite	No

Formulary ID: 24449 version 13

Last Updated: 06/20/2024 Effective: 07/01/2024



CAYSTON

Products Affected

• CAYSTON

PA Criteria	Criteria Details
Exclusion Criteria	None
Required Medical Information	Diagnosis of cystic fibrosis (confirmed by appropriate diagnostic or genetic testing) and patient has Pseudomonas aeruginosa lung infection confirmed by positive culture
Age Restrictions	7 years of age and older
Prescriber Restrictions	None
Coverage Duration	12 months
Other Criteria	None
Indications	All Medically-accepted Indications.
Off-Label Uses	N/A
Part B Prerequisite	No

Formulary ID: 24449 version 13

Last Updated: 06/20/2024 Effective: 07/01/2024



CNS STIMULANTS

Products Affected

• armodafinil

• modafinil oral

PA Criteria	Criteria Details
Exclusion Criteria	None
Required Medical Information	Diagnosis of one of the following A.) Obstructive sleep apnea (OSA) confirmed by sleep lab evaluation, B.) Narcolepsy confirmed by sleep lab evaluation, or C.) Shift work disorder (SWD)
Age Restrictions	None
Prescriber Restrictions	None
Coverage Duration	12 months
Other Criteria	None
Indications	All Medically-accepted Indications.
Off-Label Uses	N/A
Part B Prerequisite	No

Formulary ID: 24449 version 13 Last Updated: 06/20/2024

Effective: 07/01/2024



COMETRIQ

Products Affected

- COMETRIQ (100 MG DAILY DOSE) ORAL KIT 80 & 20 MG
- COMETRIQ (140 MG DAILY DOSE) ORAL KIT 3 X 20 MG & 80 MG
- COMETRIQ (60 MG DAILY DOSE)

PA Criteria	Criteria Details
Exclusion Criteria	None
Required Medical Information	Diagnosis of progressive, metastatic medullary thyroid cancer
Age Restrictions	18 years of age and older
Prescriber Restrictions	None
Coverage Duration	12 months
Other Criteria	None
Indications	All Medically-accepted Indications.
Off-Label Uses	N/A
Part B Prerequisite	No

Formulary ID: 24449 version 13 Last Updated: 06/20/2024

Effective: 07/01/2024



COPIKTRA

Products Affected

COPIKTRA

PA Criteria	Criteria Details
Exclusion Criteria	None
Required Medical Information	Diagnosis of one of the following A.) Relapsed or refractory chronic lymphocytic leukemia (CLL), or B.) Relapsed or refractory small lymphocytic lymphoma (SLL). For CLL or SLL, the patient must have history of at least 2 prior therapies
Age Restrictions	18 years of age and older
Prescriber Restrictions	Prescribed by or in consultation with an oncologist or hematologist
Coverage Duration	12 months
Other Criteria	None
Indications	All Medically-accepted Indications.
Off-Label Uses	N/A
Part B Prerequisite	No

Formulary ID: 24449 version 13

Last Updated: 06/20/2024 Effective: 07/01/2024



CORLANOR

Products Affected

• CORLANOR ORAL TABLET

PA Criteria	Criteria Details
Exclusion Criteria	Any of the following A.) Decompensated acute heart failure, B.) hypotension (i.e. blood pressure less than 90/50 mmHg), C.) sick sinus syndrome or sinoatrial block or 3rd degree AV block (unless a functioning demand pacemaker is present), D.) bradycardia (i.e., resting heart rate less than 60 bpm prior to treatment), E.) Severe hepatic impairment (Child-Pugh C), F.) Pacemaker dependent (heart rate maintained exclusively by the pacemaker), G.) Concomitant use of strong CYP3A4 inhibitors
Required Medical Information	Diagnosis of one of the following A.) Adult patients with stable, symptomatic chronic heart failure with left ventricular ejection fraction 35% or less, who are in sinus rhythm with resting heart rate 70 beats per minute or more and either are on maximally tolerated doses of betablockers or have a contraindication to beta-blocker use, or B.) Pediatric patients with stable, symptomatic heart failure due to dilated cardiomyopathy and are in sinus rhythm with an elevated heart rate
Age Restrictions	None
Prescriber Restrictions	None
Coverage Duration	12 months
Other Criteria	None
Indications	All Medically-accepted Indications.
Off-Label Uses	N/A
Part B Prerequisite	No

Formulary ID: 24449 version 13 Last Updated: 06/20/2024

Effective: 07/01/2024



COSENTYX

Products Affected

- COSENTYX (300 MG DOSE)
- COSENTYX SENSOREADY (300 MG)
- O COSENTYX SUBCUTANEOUS SOLUTION PREFILLED SYRINGE 75 MG/0.5ML
- COSENTYX UNOREADY

PA Criteria	Criteria Details
Exclusion Criteria	None
Required Medical Information	Diagnosis of one of the following A.) Ankylosing spondylitis and patient has trial and failure, contraindication, or intolerance to two preferred products, (i.e. Enbrel, Humira, Rinvoq), B.) Moderate to severe plaque psoriasis in adults and patient has trail and failure, contraindication, or intolerance to two preferred products, (i.e. Enbrel, Humira, Skyrizi, Stelara), C.) Moderate to severe plaque psoriasis in patients 6 years to less than 18 years of age and patient has trial and failure, contraindication, or intolerance to two preferred products, (i.e. Enbrel, Stelara), D.) Active psoriatic arthritis in adult patient and has trial and failure, contraindication, or intolerance to two preferred products, (i.e. Enbrel, Humira, Rinvoq, Skyrizi, Stelara), E.) Active psoriatic arthritis in patients 2 years to less than 18 years of age, F.) Non-radiographic axial spondyloarthritis and patient has trail and failure, contraindication, or intolerance to one preferred product, (i.e. Rinvoq), G.) Active enthesitis-related arthritis, or H.) Moderate to severe hidradenitis suppurativa in adults and patient has trial and failure, contraindication, or intolerance to one preferred product, (i.e. Humira)
Age Restrictions	None
Prescriber Restrictions	None
Coverage Duration	12 months
Other Criteria	Screening for latent tuberculosis infection is required prior to initiation of treatment

Formulary ID: 24449 version 13 Last Updated: 06/20/2024

Effective: 07/01/2024



PA Criteria	Criteria Details
Indications	All Medically-accepted Indications.
Off-Label Uses	N/A
Part B Prerequisite	No

Formulary ID: 24449 version 13 Last Updated: 06/20/2024

Effective: 07/01/2024



COTELLIC

Products Affected

• COTELLIC

PA Criteria	Criteria Details
Exclusion Criteria	None
Required Medical Information	Diagnosis of one of the following A.)unresectable or metastatic malignant melanoma with BRAF V600E OR V600K mutation, and documentation of combination therapy with vemurafenib (Zelboraf), or B.) Histiocytic neoplasms
Age Restrictions	None
Prescriber Restrictions	None
Coverage Duration	12 months
Other Criteria	None
Indications	All Medically-accepted Indications.
Off-Label Uses	N/A
Part B Prerequisite	No

Formulary ID: 24449 version 13

Last Updated: 06/20/2024 Effective: 07/01/2024



CYSTAGON

Products Affected

CYSTAGON

PA Criteria	Criteria Details
Exclusion Criteria	Known serious hypersensitivity to penicillamine or cysteamine
Required Medical Information	Diagnosis of nephropathic cystinosis confirmed by the presence of increased cystine concentration in leukocytes or by genetic testing
Age Restrictions	None
Prescriber Restrictions	None
Coverage Duration	12 months
Other Criteria	None
Indications	All Medically-accepted Indications.
Off-Label Uses	N/A
Part B Prerequisite	No

Formulary ID: 24449 version 13

Last Updated: 06/20/2024 Effective: 07/01/2024

 $H9306_24_DRS_09\text{-}001_OE_C$



CYSTEAMINE OPHTH

Products Affected

CYSTADROPS

CYSTARAN

PA Criteria	Criteria Details
Exclusion Criteria	None
Required Medical Information	Diagnosis of cystinosis and patient has corneal cystine crystal accumulation
Age Restrictions	None
Prescriber Restrictions	None
Coverage Duration	12 months
Other Criteria	None
Indications	All Medically-accepted Indications.
Off-Label Uses	N/A
Part B Prerequisite	No

Formulary ID: 24449 version 13

Last Updated: 06/20/2024 Effective: 07/01/2024



DALFAMPRIDINE

Products Affected

• dalfampridine er

PA Criteria	Criteria Details
Exclusion Criteria	Any of the following A.) History of seizure. B.) Moderate or severe renal impairment (creatinine clearance less than or equal to 50 mL/minute)
Required Medical Information	Diagnosis of multiple sclerosis and patient must demonstrate sustained walking impairment, but with the ability to walk 25 feet (with or without assistance) prior to starting dalfampridine
Age Restrictions	18 years of age and older
Prescriber Restrictions	Prescribed by or in consultation with a neurologist
Coverage Duration	12 months
Other Criteria	None
Indications	All Medically-accepted Indications.
Off-Label Uses	N/A
Part B Prerequisite	No

Formulary ID: 24449 version 13

Last Updated: 06/20/2024 Effective: 07/01/2024



DAURISMO

Products Affected

DAURISMO

PA Criteria	Criteria Details
Exclusion Criteria	None
Required Medical Information	Diagnosis of newly diagnosed acute myeloid leukemia (AML) and used in combination with cytarabine in patients 75 years of age or older OR in patients that have comorbidities that preclude use of intensive induction chemotherapy
Age Restrictions	18 years of age and older
Prescriber Restrictions	Prescribed by or in consultation with an oncologist or hematologist
Coverage Duration	12 months
Other Criteria	None
Indications	All Medically-accepted Indications.
Off-Label Uses	N/A
Part B Prerequisite	No

Formulary ID: 24449 version 13 Last Updated: 06/20/2024

Effective: 07/01/2024



DAYBUE

Products Affected

• DAYBUE

PA Criteria	Criteria Details
Exclusion Criteria	None
Required Medical Information	Diagnosis of Rett syndrome
Age Restrictions	2 years of age and older
Prescriber Restrictions	None
Coverage Duration	6 months
Other Criteria	None
Indications	All Medically-accepted Indications.
Off-Label Uses	N/A
Part B Prerequisite	No

Formulary ID: 24449 version 13

Last Updated: 06/20/2024 Effective: 07/01/2024



DEFERASIROX

Products Affected

• deferasirox granules

• deferasirox oral tablet

• deferasirox oral tablet soluble

PA Criteria	Criteria Details
Exclusion Criteria	Any of the following A.) Creatinine clearance less than 40 mL/min, B.) Poor performance status, C.) Platelet count less than 50 x 10(9)/L, D.) Advanced malignancy, E.) High-risk myelodysplastic syndrome (MDS)
Required Medical Information	Diagnosis of one of the following A.) Chronic iron overload in patients with non-transfusion-dependent thalassemia syndromes who have liver iron concentrations of at least 5 mg Fe/g dry weight AND serum ferritin level greater than 300 mcg/L, or B.) Chronic iron overload due to blood transfusions (transfusion hemosiderosis) as evidenced by transfusion of at least 100 mL/kg packed red blood cells AND serum ferritin level greater than 1000 mcg/L
Age Restrictions	None
Prescriber Restrictions	None
Coverage Duration	12 months
Other Criteria	None
Indications	All Medically-accepted Indications.
Off-Label Uses	N/A
Part B Prerequisite	No

Formulary ID: 24449 version 13

Last Updated: 06/20/2024 Effective: 07/01/2024



DEFERIPRONE

Products Affected

- deferiprone
- FERRIPROX ORAL SOLUTION

• FERRIPROX TWICE-A-DAY

PA Criteria	Criteria Details
Exclusion Criteria	None
Required Medical Information	Must meet all of the following 1.) Diagnosis of transfusional iron overload due to thalassemia syndromes, sickle cell disease, or other anemias, 2.) Patient has failed prior chelation therapy, and 3.) Patient has an absolute neutrophil count greater than 1.5 x 10(9)/L
Age Restrictions	3 years of age and older
Prescriber Restrictions	None
Coverage Duration	12 months
Other Criteria	None
Indications	All Medically-accepted Indications.
Off-Label Uses	N/A
Part B Prerequisite	No

Formulary ID: 24449 version 13 Last Updated: 06/20/2024

Effective: 07/01/2024



DIACOMIT

Products Affected

• DIACOMIT

PA Criteria	Criteria Details
Exclusion Criteria	None
Required Medical Information	Diagnosis of severe myoclonic epilepsy in infancy (Dravet syndrome) in patients taking clobazam
Age Restrictions	None
Prescriber Restrictions	Prescribed by or in consultation with a neurologist
Coverage Duration	12 months
Other Criteria	None
Indications	All Medically-accepted Indications.
Off-Label Uses	N/A
Part B Prerequisite	No

Formulary ID: 24449 version 13 Last Updated: 06/20/2024

Effective: 07/01/2024



DICLOFENAC TOPICAL

Products Affected

• diclofenac sodium external gel 3 %

PA Criteria	Criteria Details
Exclusion Criteria	None
Required Medical Information	Diagnosis of Actinic keratosis
Age Restrictions	None
Prescriber Restrictions	None
Coverage Duration	12 months
Other Criteria	None
Indications	All Medically-accepted Indications.
Off-Label Uses	N/A
Part B Prerequisite	No

Formulary ID: 24449 version 13 Last Updated: 06/20/2024

Effective: 07/01/2024



DIFICID

Products Affected

• DIFICID

PA Criteria	Criteria Details
Exclusion Criteria	None
Required Medical Information	Diagnosis of diarrhea associated with clostridioides difficile infection and patient has had an inadequate treatment response, intolerance, or contraindication to generic oral vancomycin
Age Restrictions	None
Prescriber Restrictions	None
Coverage Duration	4 weeks
Other Criteria	None
Indications	All Medically-accepted Indications.
Off-Label Uses	N/A
Part B Prerequisite	No

Formulary ID: 24449 version 13

Last Updated: 06/20/2024 Effective: 07/01/2024



DIMETHYL FUMARATE

Products Affected

• dimethyl fumarate oral

• dimethyl fumarate starter pack oral capsule delayed release therapy pack

PA Criteria	Criteria Details
Exclusion Criteria	None
Required Medical Information	Diagnosis of one of the following A.) Relapsing forms of multiple sclerosis (e.g., clinically isolated syndrome, relapsing-remitting MS, active secondary progressive disease, or progressive-relapsing MS), or B.) Patient has experienced a first clinical episode and has MRI features consistent with multiple sclerosis
Age Restrictions	18 years of age and older
Prescriber Restrictions	Prescribed by or in consultation with a neurologist
Coverage Duration	12 months
Other Criteria	None
Indications	All Medically-accepted Indications.
Off-Label Uses	N/A
Part B Prerequisite	No

Formulary ID: 24449 version 13

Last Updated: 06/20/2024 Effective: 07/01/2024



DOJOLVI

Products Affected

DOJOLVI

PA Criteria	Criteria Details
Exclusion Criteria	None
Required Medical Information	Diagnosis of Long-chain fatty acid oxidation disorder (LC-FAOD)
Age Restrictions	None
Prescriber Restrictions	None
Coverage Duration	12 months
Other Criteria	None
Indications	All Medically-accepted Indications.
Off-Label Uses	N/A
Part B Prerequisite	No

Formulary ID: 24449 version 13

Last Updated: 06/20/2024 Effective: 07/01/2024



DRONABINOL

Products Affected

• dronabinol

PA Criteria	Criteria Details
Exclusion Criteria	Sesame oil hypersensitivity
Required Medical Information	Diagnosis of one of the following A.) Anorexia associated to AIDS, or B.) Chemotherapy-induced nausea and vomiting
Age Restrictions	None
Prescriber Restrictions	None
Coverage Duration	6 months
Other Criteria	B vs D determination required per CMS guidance
Indications	All Medically-accepted Indications.
Off-Label Uses	N/A
Part B Prerequisite	No

Formulary ID: 24449 version 13 Last Updated: 06/20/2024

Effective: 07/01/2024



DROXIDOPA

Products Affected

• droxidopa

PA Criteria	Criteria Details
Exclusion Criteria	None
Required Medical Information	Diagnosis of symptomatic neurogenic orthostatic hypotension (nOH) caused by primary autonomic failure (e.g., Parkinson disease, multiple system atrophy, pure autonomic failure), dopamine beta-hydroxylase deficiency, or non-diabetic autonomic neuropathy
Age Restrictions	18 years of age and older
Prescriber Restrictions	None
Coverage Duration	12 months
Other Criteria	None
Indications	All Medically-accepted Indications.
Off-Label Uses	N/A
Part B Prerequisite	No

Formulary ID: 24449 version 13

Last Updated: 06/20/2024 Effective: 07/01/2024



DUPIXENT

Products Affected

DUPIXENT

PA Criteria	Criteria Details
Exclusion Criteria	None
Required Medical Information	Diagnosis of one of the following A.) Moderate to severe atopic dermatitis and if patient is 2 years or older has trial/failure, contraindication, or intolerance to two of the following 1.) Topical corticosteroid and/or 2.) Topical calcineurin inhibitor, B.) Eosinophilic phenotype or oral corticosteroid-dependent moderate to severe asthma and used as an adjunct treatment, or C.) Chronic rhinosinusitis with nasal polyposis and used as an adjunct treatment, D.) Eosinophilic esophagitis, or E.) Prurigo nodularis
Age Restrictions	None
Prescriber Restrictions	None
Coverage Duration	12 months
Other Criteria	None
Indications	All Medically-accepted Indications.
Off-Label Uses	N/A
Part B Prerequisite	No

Formulary ID: 24449 version 13

Last Updated: 06/20/2024 Effective: 07/01/2024

 $H9306_24_DRS_09\text{-}001_OE_C$



EMGALITY

Products Affected

• EMGALITY

PA Criteria	Criteria Details
Exclusion Criteria	None
Required Medical Information	Diagnosis of one of the following A.) Chronic or episodic migraine disorder and patient has documented trial, inadequate response, or contraindication to at least 2 generic formulary drugs used for migraine prevention (i.e., propranolol, topiramate, divalproex, timolol), or B.) Episodic cluster headache
Age Restrictions	18 years of age and older
Prescriber Restrictions	None
Coverage Duration	12 months
Other Criteria	None
Indications	All Medically-accepted Indications.
Off-Label Uses	N/A
Part B Prerequisite	No

Formulary ID: 24449 version 13

Last Updated: 06/20/2024 Effective: 07/01/2024

 $H9306_24_DRS_09\text{-}001_OE_C$



ENBREL

Products Affected

- ENBREL MINI
- ENBREL SUBCUTANEOUS SOLUTION 25 MG/0.5ML
- ENBREL SUBCUTANEOUS SOLUTION PREFILLED SYRINGE
- ENBREL SURECLICK SUBCUTANEOUS SOLUTION AUTO-INJECTOR

SOLUTION I REFIELED STRINGE	
PA Criteria	Criteria Details
Exclusion Criteria	None
Required Medical Information	Diagnosis of one of the following A.) Moderate to severe rheumatoid arthritis, B.) Moderate to severe polyarticular juvenile idiopathic arthritis, C.) Psoriatic arthritis, D.) Ankylosing spondylitis, or E.) Moderate to severe chronic plaque psoriasis in patients who are candidates for systemic therapy or phototherapy
Age Restrictions	None
Prescriber Restrictions	None
Coverage Duration	12 months
Other Criteria	Screening for latent tuberculosis infection is required prior to initiation of treatment
Indications	All Medically-accepted Indications.
Off-Label Uses	N/A
Part B Prerequisite	No

Formulary ID: 24449 version 13 Last Updated: 06/20/2024

Effective: 07/01/2024



ENDARI

Products Affected

ENDARI

PA Criteria	Criteria Details
Exclusion Criteria	None
Required Medical Information	Diagnosis of sickle cell disease AND one of the following 1.) Patient has acute complications and is being treated with Hydroxyurea, or 2.) Patient has acute complications and is unable to tolerate Hydroxyurea
Age Restrictions	5 years of age and older
Prescriber Restrictions	None
Coverage Duration	12 months
Other Criteria	None
Indications	All Medically-accepted Indications.
Off-Label Uses	N/A
Part B Prerequisite	No

Formulary ID: 24449 version 13

Last Updated: 06/20/2024 Effective: 07/01/2024



ENSPRYNG

Products Affected

ENSPRYNG

PA Criteria	Criteria Details
Exclusion Criteria	Any of the following A.) Active Hepatitis B infection, or B.) Active or untreated latent tuberculosis
Required Medical Information	Diagnosis of neuromyelitis optica spectrum disorder (NMOSD) in patients who are anti-aquaporin-4 (AQP4) antibody positive
Age Restrictions	18 years of age and older
Prescriber Restrictions	Prescribed by or in consultation with a neurologist, immunologist, or ophthalmologist
Coverage Duration	12 months
Other Criteria	None
Indications	All Medically-accepted Indications.
Off-Label Uses	N/A
Part B Prerequisite	No

Formulary ID: 24449 version 13

Last Updated: 06/20/2024 Effective: 07/01/2024



EPIDIOLEX

Products Affected

• EPIDIOLEX

PA Criteria	Criteria Details
Exclusion Criteria	None
Required Medical Information	Diagnosis of one of the following A.) Lennox-Gastaut syndrome, B.) Severe myoclonic epilepsy in infancy (Dravet syndrome), or C.) Seizures associated with tuberous sclerosis complex (TSC)
Age Restrictions	None
Prescriber Restrictions	Prescribed by or in consultation with a neurologist
Coverage Duration	12 months
Other Criteria	None
Indications	All Medically-accepted Indications.
Off-Label Uses	N/A
Part B Prerequisite	No

Formulary ID: 24449 version 13

Last Updated: 06/20/2024 Effective: 07/01/2024



EPOETIN THERAPY

Products Affected

• RETACRIT INJECTION SOLUTION 10000 UNIT/ML, 10000 UNIT/ML(1ML), 2000 UNIT/ML, 3000

UNIT/ML, 4000 UNIT/ML, 40000 UNIT/ML

PA Criteria	Criteria Details
Exclusion Criteria	None
Required Medical Information	Diagnosis of one of the following A.) Non-myeloid neoplastic disease and utilized for the treatment of chemotherapy induced anemia, B.) HIV infection and utilized for the treatment of zidovudine induced anemia, C.) Chronic kidney disease resulting in anemia, or D.) High risk surgical candidate status at risk for perioperative blood loss and undergoing elective, noncardiac, or nonvascular surgery
Age Restrictions	None
Prescriber Restrictions	None
Coverage Duration	12 months
Other Criteria	B vs D determination required per CMS guidance
Indications	All Medically-accepted Indications.
Off-Label Uses	N/A
Part B Prerequisite	No

Formulary ID: 24449 version 13 Last Updated: 06/20/2024

Effective: 07/01/2024



ERIVEDGE

Products Affected

• ERIVEDGE

PA Criteria	Criteria Details
Exclusion Criteria	Pregnancy
Required Medical Information	Diagnosis of one of the following A.) Metastatic basal cell carcinoma, or B.) Locally advanced basal cell carcinoma that has recurred following surgery or the patient is not a candidate for surgery or radiation
Age Restrictions	18 years of age and older
Prescriber Restrictions	None
Coverage Duration	12 months
Other Criteria	None
Indications	All Medically-accepted Indications.
Off-Label Uses	N/A
Part B Prerequisite	No

Formulary ID: 24449 version 13 Last Updated: 06/20/2024

Effective: 07/01/2024



ERLEADA

Products Affected

ERLEADA

PA Criteria	Criteria Details
Exclusion Criteria	None
Required Medical Information	Diagnosis of one of the following A.) Nonmetastatic, castration-resistant prostate cancer (nmCRPC), or B.) Metastatic, castration-sensitive prostate cancer (mCSPC). For treatment of nmCRPC and mCSPC, one of the following applies 1.) Used in combination with a gonadotropin-releasing hormone (GnRH) analog, OR 2) Patient has received bilateral orchiectomy
Age Restrictions	18 years of age and older
Prescriber Restrictions	Prescribed by or in consultation with an oncologist or urologist
Coverage Duration	12 months
Other Criteria	None
Indications	All Medically-accepted Indications.
Off-Label Uses	N/A
Part B Prerequisite	No

Formulary ID: 24449 version 13 Last Updated: 06/20/2024

Effective: 07/01/2024



ERLOTINIB

Products Affected

• erlotinib hcl

PA Criteria	Criteria Details
Exclusion Criteria	None
Required Medical Information	Diagnosis of one of the following A.) Locally advanced, unresectable, or metastatic pancreatic cancer and erlotinib will be used in combination with gemcitabine, B.) Locally advanced or metastatic non-small cell lung cancer with epidermal growth factor receptor (EGFR) exon 19 deletions or exon 21 (L858R) substitution mutations as detected by an FDA-approved test or Clinical Laboratory Improvement Amendments-approved facility AND one of the following 1.) Erlotinib will be used as first-line treatment, 2.) Failure with at least one prior chemotherapy regimen, or 3.) No evidence of disease progression after four cycles of first-line platinum-based chemotherapy and erlotinib will be used as maintenance treatment
Age Restrictions	18 years of age and older
Prescriber Restrictions	Prescribed by or in consultation with an oncologist
Coverage Duration	12 months
Other Criteria	None
Indications	All Medically-accepted Indications.
Off-Label Uses	N/A
Part B Prerequisite	No

Formulary ID: 24449 version 13

Last Updated: 06/20/2024 Effective: 07/01/2024



EVEROLIMUS

Products Affected

• everolimus oral tablet 10 mg, 2.5 mg, 5 mg, 7.5 mg

PA Criteria	Criteria Details
Exclusion Criteria	Any of the following A.) Hypersensitivity to everolimus or excipients, or B.) Hypersensitivity to rapamycin derivatives (e.g. sirolimus)
Required Medical Information	Diagnosis of one of the following A.) Renal angiomyolipoma and tuberous sclerosis complex (TSC) not requiring immediate surgery, B.) Advanced hormone receptor-positive, HER2 negative breast cancer in postmenopausal women and taken in combination with exemestane, after failure with letrozole or anastrozole, C.) Progressive, well-differentiated, nonfunctional neuroendocrine tumors of gastrointestinal or lung origin and disease is unresectable, locally advanced, or metastatic, D.) Pancreatic progressive neuroendocrine tumors and disease is unresectable, locally advanced, or metastatic, E.) Advanced renal cell carcinoma (RCC) after failure with sunitinib or sorafenib, F.) Subependymal giant cell astrocytoma (SEGA) associated with tuberous sclerosis complex in patients who are not candidates for curative surgical resection
Age Restrictions	None
Prescriber Restrictions	Prescribed by or in consultation with an oncologist or neurologist
Coverage Duration	12 months
Other Criteria	None
Indications	All Medically-accepted Indications.
Off-Label Uses	N/A
Part B Prerequisite	No

Formulary ID: 24449 version 13 Last Updated: 06/20/2024

Effective: 07/01/2024



EVEROLIMUS SUSPENSION

Products Affected

• everolimus oral tablet soluble

PA Criteria	Criteria Details
Exclusion Criteria	Any of the following A.) Hypersensitivity to everolimus, or B.) Hypersensitivity to rapamycin derivatives (e.g. sirolimus)
Required Medical Information	Diagnosis of one of the following A.) Tuberous sclerosis complex (TSC)-associated partial-onset seizures, or B.) Subependymal giant cell astrocytoma (SEGA) associated with tuberous sclerosis complex in patients who are not candidates for curative surgical resection
Age Restrictions	None
Prescriber Restrictions	Prescribed by or in consultation with an oncologist or neurologist
Coverage Duration	12 months
Other Criteria	None
Indications	All Medically-accepted Indications.
Off-Label Uses	N/A
Part B Prerequisite	No

Formulary ID: 24449 version 13

Last Updated: 06/20/2024 Effective: 07/01/2024



EVRYSDI

Products Affected

EVRYSDI

PA Criteria	Criteria Details
Exclusion Criteria	None
Required Medical Information	Diagnosis of spinal muscular atrophy (SMA)
Age Restrictions	None
Prescriber Restrictions	Prescribed by or in consultation with a neurologist
Coverage Duration	12 months
Other Criteria	None
Indications	All Medically-accepted Indications.
Off-Label Uses	N/A
Part B Prerequisite	No

Formulary ID: 24449 version 13

Last Updated: 06/20/2024 Effective: 07/01/2024

 $H9306_24_DRS_09\text{-}001_OE_C$



EXKIVITY

Products Affected

EXKIVITY

PA Criteria	Criteria Details
Exclusion Criteria	None
Required Medical Information	Diagnosis of locally advanced or metastatic non-small cell lung cancer (NSCLC) with EGFR exon 20 insertion mutations (as confirmed by an FDA-approved test) AND whose disease has progressed on or after platinum-based chemotherapy
Age Restrictions	18 years of age and older
Prescriber Restrictions	Prescribed by or in consultation with an oncologist
Coverage Duration	12 months
Other Criteria	None
Indications	All Medically-accepted Indications.
Off-Label Uses	N/A
Part B Prerequisite	No

Formulary ID: 24449 version 13

Last Updated: 06/20/2024 Effective: 07/01/2024

 $H9306_24_DRS_09\text{-}001_OE_C$



FEBUXOSTAT

Products Affected

• febuxostat

PA Criteria	Criteria Details
Exclusion Criteria	Concomitant use of azathioprine or mercaptopurine
Required Medical Information	Diagnosis of Gout and all of the following 1.) documented inadequate treatment response, adverse event, or contraindication to maximally titrated dose of Allopurinol, and 2.) patients with established cardiovascular disease, prescriber attests that benefit of treatment outweighs the risk of treatment
Age Restrictions	18 years of age and older
Prescriber Restrictions	None
Coverage Duration	12 months
Other Criteria	None
Indications	All Medically-accepted Indications.
Off-Label Uses	N/A
Part B Prerequisite	No

Formulary ID: 24449 version 13 Last Updated: 06/20/2024

Effective: 07/01/2024



FENTANYL ORAL

Products Affected

• fentanyl citrate buccal lozenge on a handle

PA Criteria	Criteria Details
Exclusion Criteria	Any of the following A.) Management of acute or postoperative pain (including headache/migraine, dental pain, and use in the emergency room), B.) Use in opioid non-tolerant patients, C.) Known or suspected gastrointestinal obstruction, including paralytic ileus, D.) Acute or severe bronchial asthma and used in an unmonitored setting (absence of resuscitative equipment)
Required Medical Information	Must meet all of the following 1.) Diagnosis of cancer-related breakthrough pain, 2.) Patient is currently receiving/tolerant to around-the-clock opioid therapy for persistent cancer pain, and 3.) Patient and prescriber are enrolled in the TIRF REMS Access Program
Age Restrictions	16 years of age and older
Prescriber Restrictions	None
Coverage Duration	12 months
Other Criteria	None
Indications	All Medically-accepted Indications.
Off-Label Uses	N/A
Part B Prerequisite	No

Formulary ID: 24449 version 13 Last Updated: 06/20/2024

Effective: 07/01/2024



FENTANYL PATCH

Products Affected

• fentanyl

PA Criteria	Criteria Details
Exclusion Criteria	Any of the following A.) Management of acute or postoperative pain (including headache/migraine, dental pain, and use in the emergency room), B.) Mild or intermittent pain management, C.) Use in opioid nontolerant patients, D.) Known or suspected gastrointestinal obstruction, including paralytic ileus, E.) Acute or severe bronchial asthma and used in an unmonitored setting (absence of resuscitative equipment)
Required Medical Information	Must meet all of the following 1.) Patient is opioid tolerant (taking for one week or longer at least 60mg of morphine or equivalent daily) and 2.) Patient has tried at least one extended release oral opioids or is unable to take extended release oral opioids secondary to allergy, adverse events, swallowing difficulty, or uncontrollable nausea/vomiting
Age Restrictions	None
Prescriber Restrictions	None
Coverage Duration	12 months
Other Criteria	None
Indications	All Medically-accepted Indications.
Off-Label Uses	N/A
Part B Prerequisite	No

Formulary ID: 24449 version 13 Last Updated: 06/20/2024

Effective: 07/01/2024



FILSPARI

Products Affected

• FILSPARI

PA Criteria	Criteria Details
Exclusion Criteria	Any of the following A.) Pregnancy or B.) Concomitant use with angiotensin receptor blockers (ARBs), endothelin receptor antagonists (ERAs), or aliskiren
Required Medical Information	Diagnosis of treatment of primary immunoglobulin A (IgA) nephropathy at risk of rapid disease progression, generally a urine protein to creatinine ratio (UPCR) of 1.5 g/g or more, to reduce proteinuria
Age Restrictions	18 years of age and older
Prescriber Restrictions	None
Coverage Duration	12 months
Other Criteria	None
Indications	All Medically-accepted Indications.
Off-Label Uses	N/A
Part B Prerequisite	No

Formulary ID: 24449 version 13 Last Updated: 06/20/2024

Effective: 07/01/2024



FINGOLIMOD

Products Affected

• fingolimod hcl

PA Criteria	Criteria Details
Exclusion Criteria	Any of the following A.) Recent (within the last 6 months) occurrence of: myocardial infarction, unstable angina, stroke, transient ischemic attack, decompensated heart failure requiring hospitalization, or Class III/IV heart failure, B.) History or presence of Mobitz Type II 2nd degree or 3rd degree AV block or sick sinus syndrome, unless patient has a pacemaker, C.) Baseline QTC interval greater than or equal to 500 milliseconds, D.) Receiving concurrent treatment with Class Ia or Class III anti-arrhythmic drugs (quinidine, procainamide, amiodarone, sotalol)
Required Medical Information	Diagnosis of one of the following A.) Relapsing forms of multiple sclerosis (e.g., clinically isolated syndrome, relapsing-remitting MS, active secondary progressive disease, or progressive-relapsing MS), or B.) Patient has experienced a first clinical episode and has MRI features consistent with multiple sclerosis
Age Restrictions	10 years of age and older
Prescriber Restrictions	Prescribed by or in consultation with a neurologist
Coverage Duration	12 months
Other Criteria	None
Indications	All Medically-accepted Indications.
Off-Label Uses	N/A
Part B Prerequisite	No

Formulary ID: 24449 version 13 Last Updated: 06/20/2024

Effective: 07/01/2024



FINTEPLA

Products Affected

• FINTEPLA

PA Criteria	Criteria Details
Exclusion Criteria	Any of the following A.) Concomitant use of an MAOI, or B.) Use within 14 days of discontinuing an MAOI
Required Medical Information	Diagnosis of Severe myoclonic epilepsy in infancy (Dravet syndrome) or seizures associated with Lennox-Gastaut syndrome
Age Restrictions	None
Prescriber Restrictions	Prescribed by or in consultation with a neurologist
Coverage Duration	12 months
Other Criteria	None
Indications	All Medically-accepted Indications.
Off-Label Uses	N/A
Part B Prerequisite	No

Formulary ID: 24449 version 13 Last Updated: 06/20/2024

Effective: 07/01/2024



FIRMAGON

Products Affected

- FIRMAGON (240 MG DOSE)
- FIRMAGON SUBCUTANEOUS SOLUTION RECONSTITUTED 80 MG

PA Criteria	Criteria Details
Exclusion Criteria	none
Required Medical Information	Diagnosis of advanced prostate cancer
Age Restrictions	none
Prescriber Restrictions	none
Coverage Duration	12 months
Other Criteria	B vs D determination required per CMS guidance
Indications	All Medically-accepted Indications.
Off-Label Uses	N/A
Part B Prerequisite	No

Formulary ID: 24449 version 13 Last Updated: 06/20/2024

Effective: 07/01/2024



FOTIVDA

Products Affected

FOTIVDA

PA Criteria	Criteria Details
Exclusion Criteria	None
Required Medical Information	Diagnosis of relapsed or refractory advanced renal cell cancer (RCC) following 2 or more prior systemic therapies
Age Restrictions	18 years of age and older
Prescriber Restrictions	Prescribed by or in consultation with an oncologist
Coverage Duration	12 months
Other Criteria	None
Indications	All Medically-accepted Indications.
Off-Label Uses	N/A
Part B Prerequisite	No

Formulary ID: 24449 version 13 Last Updated: 06/20/2024

Effective: 07/01/2024



FRUZAQLA

Products Affected

• FRUZAQLA

PA Criteria	Criteria Details
Exclusion Criteria	None
Required Medical Information	Diagnosis of metastatic colorectal cancer (mCRC) and all of the following: A.) patient has been previously treated with fluoropyrimidine, oxaliplatin, irinotecan-based chemotherapy, B.) an anti-VEGF therapy, and C.) if RAS wild-type and medically appropriate, patient has also been previously treated with anti-EGFR therapy
Age Restrictions	18 years of age and older
Prescriber Restrictions	Prescribed by or in consultation with an oncologist or hematologist
Coverage Duration	3 months
Other Criteria	None
Indications	All Medically-accepted Indications.
Off-Label Uses	N/A
Part B Prerequisite	No

Formulary ID: 24449 version 13 Last Updated: 06/20/2024

Effective: 07/01/2024



GALAFOLD

Products Affected

GALAFOLD

PA Criteria	Criteria Details
Exclusion Criteria	None
Required Medical Information	Diagnosis of Fabry disease with an amenable galactosidase alpha gene (GLA) mutation
Age Restrictions	16 years of age and older
Prescriber Restrictions	None
Coverage Duration	12 months
Other Criteria	None
Indications	All Medically-accepted Indications.
Off-Label Uses	N/A
Part B Prerequisite	No

Formulary ID: 24449 version 13 Last Updated: 06/20/2024

Effective: 07/01/2024



GATTEX

Products Affected

• GATTEX

PA Criteria	Criteria Details
Exclusion Criteria	None
Required Medical Information	Diagnosis of short bowel syndrome and patient is dependent on parenteral support
Age Restrictions	None
Prescriber Restrictions	None
Coverage Duration	12 months
Other Criteria	None
Indications	All Medically-accepted Indications.
Off-Label Uses	N/A
Part B Prerequisite	No

Formulary ID: 24449 version 13

Last Updated: 06/20/2024 Effective: 07/01/2024

 $H9306_24_DRS_09\text{-}001_OE_C$



GAVRETO

Products Affected

GAVRETO

PA Criteria	Criteria Details
Exclusion Criteria	None
Required Medical Information	Diagnosis of one of the following A.) Metastatic RET fusion-positive non-small cell lung cancer (NSCLC) as detected by an FDA approved test, B.) Advanced or metastatic RET-mutant medullary thyroid cancer and patient requires systemic therapy, or C.) Advanced or metastatic RET fusion-positive thyroid and patient requires systemic therapy and is radioactive iodine-refractory, when radioactive iodine is appropriate
Age Restrictions	None
Prescriber Restrictions	Prescribed by or in consultation with an oncologist
Coverage Duration	12 months
Other Criteria	None
Indications	All Medically-accepted Indications.
Off-Label Uses	N/A
Part B Prerequisite	No

Formulary ID: 24449 version 13 Last Updated: 06/20/2024

Effective: 07/01/2024



GEFITINIB

Products Affected

• gefitinib

PA Criteria	Criteria Details
Exclusion Criteria	None
Required Medical Information	Diagnosis of metastatic non-small cell lung cancer (NSCLC) and must meet all of the following 1.) Tumor has epidermal growth factor receptor (EGFR) exon 19 deletions or exon 21 (L858R) substitution mutations as detected by an FDA-approved test or Clinical Laboratory Improvement Amendments-approved facility and 2.) Used as first-line treatment
Age Restrictions	18 years of age and older
Prescriber Restrictions	None
Coverage Duration	12 months
Other Criteria	None
Indications	All Medically-accepted Indications.
Off-Label Uses	N/A
Part B Prerequisite	No

Formulary ID: 24449 version 13

Last Updated: 06/20/2024 Effective: 07/01/2024



GILOTRIF

Products Affected

• GILOTRIF

PA Criteria	Criteria Details
Exclusion Criteria	None
Required Medical Information	Diagnosis of one of the following A.) Metastatic non-small cell lung cancer (NSCLC) in patients whose tumors have non-resistant epidermal growth factor receptor (EGFR) mutations as detected by an FDA-approved test, or B.) Metastatic squamous NSCLC with progression after platinum-based chemotherapy
Age Restrictions	18 years of age and older
Prescriber Restrictions	Prescribed by or in consultation with an oncologist
Coverage Duration	12 months
Other Criteria	None
Indications	All Medically-accepted Indications.
Off-Label Uses	N/A
Part B Prerequisite	No

Formulary ID: 24449 version 13 Last Updated: 06/20/2024

Effective: 07/01/2024



GLATIRAMER

Products Affected

• COPAXONE SUBCUTANEOUS SOLUTION PREFILLED SYRINGE

PA Criteria	Criteria Details
Exclusion Criteria	None
Required Medical Information	Diagnosis of one of the following A.) Relapsing forms of multiple sclerosis (e.g., clinically isolated syndrome, relapsing-remitting MS, active secondary progressive disease, or progressive-relapsing MS), or B.) Patient has experienced a first clinical episode and has MRI features consistent with multiple sclerosis
Age Restrictions	18 years of age and older
Prescriber Restrictions	Prescribed by or in consultation with a neurologist
Coverage Duration	12 months
Other Criteria	None
Indications	All Medically-accepted Indications.
Off-Label Uses	N/A
Part B Prerequisite	No

Formulary ID: 24449 version 13 Last Updated: 06/20/2024

Effective: 07/01/2024



GLEOSTINE

Products Affected

• GLEOSTINE ORAL CAPSULE 10 MG, 100 MG, 40 MG

PA Criteria	Criteria Details
Exclusion Criteria	None
Required Medical Information	Must meet one of the following: A.) Hodgkin's disease in patient who has relapsed during or failed to respond to primary therapy and is being used in combination with other agents OR B.) Intracranial tumor
Age Restrictions	None
Prescriber Restrictions	Prescribed by or in consultation with an oncologist
Coverage Duration	12 months
Other Criteria	None
Indications	All Medically-accepted Indications.
Off-Label Uses	N/A
Part B Prerequisite	No

Formulary ID: 24449 version 13 Last Updated: 06/20/2024

Effective: 07/01/2024



GLP1

Products Affected

- MOUNJARO
- OZEMPIC (0.25 OR 0.5 MG/DOSE) SUBCUTANEOUS SOLUTION PEN-INJECTOR 2 MG/3ML
- OZEMPIC (1 MG/DOSE)
 SUBCUTANEOUS SOLUTION PEN-INJECTOR 4 MG/3ML
- OZEMPIC (2 MG/DOSE)
- RYBELSUS
- TRULICITY
- VICTOZA SUBCUTANEOUS SOLUTION PEN-INJECTOR

INJECTOR 4 MIG/SMIL	
PA Criteria	Criteria Details
Exclusion Criteria	None
Required Medical Information	Must meet all of the following 1.) The drug is prescribed for an FDA-approved indication, 2.) For a diagnosis of Type 2 Diabetes Mellitus the patient has a trial and failure, contraindication or intolerance to metformin or any metformin combination product
Age Restrictions	None
Prescriber Restrictions	None
Coverage Duration	12 monthd
Other Criteria	None
Indications	All Medically-accepted Indications.
Off-Label Uses	N/A
Part B Prerequisite	No

Formulary ID: 24449 version 13 Last Updated: 06/20/2024

Effective: 07/01/2024



GROWTH HORMONE

Products Affected

- OMNITROPE SUBCUTANEOUS SOLUTION CARTRIDGE
- OMNITROPE SUBCUTANEOUS SOLUTION RECONSTITUTED

PA Criteria	Criteria Details
Exclusion Criteria	Any of the following A.) Use for growth promotion in pediatric patients with closed epiphyses, B.) Acute critical illness caused by complications following open-heart or abdominal surgery, multiple accidental trauma, or acute respiratory failure, C.) Active malignancy, D.) Active proliferative or severe nonproliferative diabetic retinopathy, E.) Prader-Willi Syndrome in patients who are severely obese, have a history of upper airway obstruction or sleep apnea, or have severe respiratory impairment
Required Medical Information	Diagnosis of pediatric indication: A.) GHD and bone age at least 1 year or 2 standard deviations (SD) delayed compared with chronological age and 2 stim tests with peak GH secretion below 10 ng/mL or IGF-1/IGFBP3 level more than 2 SDS below mean if CNS pathology, h/o irradiation, or proven genetic cause, B.) SGA and birth weight or length 2 or more SDS below mean for gestational age and fails to manifest catch up growth by age 2 (height 2 or more SDS below mean for age and gender), C.) CRI and metabolic abnormalities have been corrected, and patient has not had renal transplant D.) SHOX deficiency or Noonan syndrome E.) PWS confirmed by genetic testing, F.) Turner Syndrome confirmed by chromosome analysis. For GHD, CRI, SHOX deficiency, Noonan syndrome, and PWS one of the following height more than 3 SDS below mean for age and gender, or height more than 2 SDS below mean with GV more than 1 SDS below mean, or GV over 1 year 2 SDS below mean. OR Diagnosis of an adult indication: A.) childhood- or adult-onset GHD confirmed by 2 standard GH stim tests (provide assay): 1 test must be insulin tolerance test (ITT) with blood glucose nadir less than 40 mg/dL (2.2 mmol/L). If contraindicated, use a standardized stim test (i.e. arginine plus GH releasing hormone [preferred], glucagon, arginine), B.) GHD with at least 1 other pituitary hormone deficiency and failed at least 1 GH stim test (ITT preferred), C.) GHD with panhypopituitarism (3 or more pituitary hormone deficiencies), D.) GHD with irreversible hypothalamic-pituitary structural lesions due to tumors, surgery or radiation of pituitary or hypothalamus region AND a subnormal IGF-1 (after at least 1 month off GH therapy)

Formulary ID: 24449 version 13 Last Updated: 06/20/2024

Effective: 07/01/2024



PA Criteria	Criteria Details
	AND Objective evidence of GHD complications, such as: low bone density, increased visceral fat mass, or cardiovascular complications AND Completed linear growth (GV less than 2 cm/year) AND GH has been discontinued for at least 1 month (if previously receiving GH)
Age Restrictions	None
Prescriber Restrictions	Prescribed by or in consultation with an endocrinologist or nephrologist
Coverage Duration	12 months
Other Criteria	None
Indications	All Medically-accepted Indications.
Off-Label Uses	N/A
Part B Prerequisite	No

Formulary ID: 24449 version 13

Last Updated: 06/20/2024 Effective: 07/01/2024



HEPATITIS C

Products Affected

MAVYRET

VOSEVI

• sofosbuvir-velpatasvir

PA Criteria	Criteria Details
Exclusion Criteria	None
Required Medical Information	Diagnosis of HCV genotype, subtype and quantitative HCV RNA (viral load) testing any time prior to therapy. Must document cirrhosis status, prior treatment history (if any), and planned duration of treatment. All genotypes will require trial/failure, contraindication to, or intolerance to Mavyret or Sofosbuvir-Velpatasvir prior to the approval of Vosevi. Genotype and subtype are not required for: (1) initial treatment of patients without cirrhosis if using Sofosbuvir-Velpatasvir or Mavyret OR (2) treatment of patients with compensated cirrhosis if using Mavyret
Age Restrictions	None
Prescriber Restrictions	Prescribed by or in consultation with a gastroenterologist, hepatologist, or infectious disease specialist
Coverage Duration	Duration of approval per AASLD Guidelines
Other Criteria	None
Indications	All Medically-accepted Indications.
Off-Label Uses	N/A
Part B Prerequisite	No

Formulary ID: 24449 version 13 Last Updated: 06/20/2024

Effective: 07/01/2024



HUMIRA

Products Affected

- HUMIRA (2 PEN)
- HUMIRA (2 SYRINGE) SUBCUTANEOUS PREFILLED SYRINGE KIT 10 MG/0.1ML, 20 MG/0.2ML, 40 MG/0.4ML, 40 MG/0.8ML
- HUMIRA-CD/UC/HS STARTER SUBCUTANEOUS PEN-INJECTOR KIT 80 MG/0.8ML
- HUMIRA-PED<40KG CROHNS STARTER
- HUMIRA-PED>/=40KG CROHNS START
- HUMIRA-PED>/=40KG UC STARTER
- HUMIRA-PS/UV/ADOL HS STARTER
- HUMIRA-PSORIASIS/UVEIT STARTER

PA Criteria	Criteria Details
Exclusion Criteria	None
Required Medical Information	Diagnosis of one of the following A.) Moderate to severe rheumatoid arthritis, B.) Moderate to severe polyarticular juvenile idiopathic arthritis, C.) Psoriatic arthritis, D.) Ankylosing spondylitis, E.) Moderate to severe chronic plaque psoriasis in patients who are candidates for systemic therapy or phototherapy and when other systemic therapies are medically less appropriate, F.) Moderate to severe Crohn's disease in patients who have had an inadequate response to conventional therapy, G.) Moderate to severe ulcerative colitis in patients who have had an inadequate response to immunosuppressants (e.g. corticosteroids, azathioprine), H.) Noninfectious uveitis (including intermediate, posterior, and panuveitis), or I.) Moderate to severe hidradenitis suppurativa
Age Restrictions	None
Prescriber Restrictions	None
Coverage Duration	12 months
Other Criteria	Screening for latent tuberculosis infection is required prior to initiation of treatment
Indications	All Medically-accepted Indications.

Formulary ID: 24449 version 13 Last Updated: 06/20/2024

Effective: 07/01/2024



PA Criteria	Criteria Details
Off-Label Uses	N/A
Part B Prerequisite	No

Formulary ID: 24449 version 13 Last Updated: 06/20/2024

Effective: 07/01/2024



HYFTOR

Products Affected

• HYFTOR

PA Criteria	Criteria Details
Exclusion Criteria	None
Required Medical Information	Diagnosis of Facial angiofibroma associated with tuberous sclerosis
Age Restrictions	6 years of age and older
Prescriber Restrictions	None
Coverage Duration	12 months
Other Criteria	None
Indications	All Medically-accepted Indications.
Off-Label Uses	N/A
Part B Prerequisite	No

Formulary ID: 24449 version 13

Last Updated: 06/20/2024 Effective: 07/01/2024

 $H9306_24_DRS_09\text{-}001_OE_C$



IBRANCE

Products Affected

• IBRANCE

PA Criteria	Criteria Details
Exclusion Criteria	None
Required Medical Information	Diagnosis of one of the following A.) Advanced or metastatic, hormone receptor (HR)-positive, human epidermal growth factor receptor 2 (HER2)-negative breast cancer used in combination with fulvestrant and disease has progressed following endocrine therapy, or B.) Advanced or metastatic, hormone receptor (HR)-positive, human epidermal growth factor receptor 2 (HER2)-negative breast cancer and used in combination with an aromatase inhibitor in a male or female patient as initial endocrine-based therapy
Age Restrictions	18 years of age and older
Prescriber Restrictions	Prescribed by or in consultation with an oncologist
Coverage Duration	12 months
Other Criteria	None
Indications	All Medically-accepted Indications.
Off-Label Uses	N/A
Part B Prerequisite	No

Formulary ID: 24449 version 13

Last Updated: 06/20/2024 Effective: 07/01/2024



ICATIBANT

Products Affected

• icatibant acetate subcutaneous solution prefilled syringe

PA Criteria	Criteria Details
Exclusion Criteria	None
Required Medical Information	Diagnosis of one of the following and used as treatment for acute attacks A.) Hereditary angioedema (HAE) with C1 inhibitor deficiency (Type 1) confirmed by laboratory testing, or B.) HAE with C1 inhibitor dysfunction (Type 2) confirmed by laboratory testing, or C.) HAE with normal C1 inhibitor (Type 3) confirmed by laboratory testing and one of the following 1.) Positive test for an F12, angiopoietin-1, or plasminogen gene mutation, or 2.) Family history of angioedema and the angioedema was refractory to a trial of an antihistamine for at least one month
Age Restrictions	18 years of age and older
Prescriber Restrictions	Prescribed by or in consultation with an allergist, hematologist, or immunologist
Coverage Duration	12 months
Other Criteria	None
Indications	All Medically-accepted Indications.
Off-Label Uses	N/A
Part B Prerequisite	No

Formulary ID: 24449 version 13 Last Updated: 06/20/2024

Effective: 07/01/2024



ICLUSIG

Products Affected

• ICLUSIG

PA Criteria	Criteria Details
Exclusion Criteria	None
Required Medical Information	Diagnosis of one of the following A.) Chronic phase, accelerated phase, or blast phase chronic myeloid leukemia (CML) in adult patients who are T315I-positive or for whom no other tyrosine kinase inhibitor therapy is indicated, B.) Chronic phase, chronic myeloid leukemia (CML) in adult patients with resistance or intolerance to at least two prior kinase inhibitors, or C.) Philadelphia chromosome-positive acute lymphoblastic leukemia (Ph+ ALL) in adult patients who are T315I-positive or for whom no other tyrosine kinase inhibitor therapy is indicated
Age Restrictions	18 years of age and older
Prescriber Restrictions	Prescribed by or in consultation with an oncologist or hematologist
Coverage Duration	12 months
Other Criteria	None
Indications	All Medically-accepted Indications.
Off-Label Uses	N/A
Part B Prerequisite	No

Formulary ID: 24449 version 13 Last Updated: 06/20/2024

Effective: 07/01/2024



IDHIFA

Products Affected

• IDHIFA

PA Criteria	Criteria Details
Exclusion Criteria	None
Required Medical Information	Diagnosis of relapsed or refractory acute myeloid leukemia (AML) with an isocitrate dehydrogenase 2 (IDH2) mutation as detected by an FDA approved test
Age Restrictions	18 years of age and older
Prescriber Restrictions	Prescribed by or in consultation with an oncologist or hematologist
Coverage Duration	12 months
Other Criteria	None
Indications	All Medically-accepted Indications.
Off-Label Uses	N/A
Part B Prerequisite	No

Formulary ID: 24449 version 13

Last Updated: 06/20/2024 Effective: 07/01/2024



IMATINIB

Products Affected

• imatinib mesylate

PA Criteria	Criteria Details
Exclusion Criteria	None
Required Medical Information	Diagnosis of one of the following A.) Philadelphia chromosome-positive chronic myelogenous leukemia (Ph+ CML), B.) Ph+ acute lymphoblastic leukemia (ALL), C.) Gastrointestinal stromal tumor (GIST) where patient has documented c-KIT (CD117) positive unresectable or metastatic malignant GIST or patient had resection of c-KIT positive GIST and imatinib will be used as an adjuvant therapy, D.) Dermatofibrosarcoma protuberans that is unresectable, recurrent, or metastatic, E.) Hypereosinophilic syndrome or chronic eosinophilic leukemia, F.) Myelodysplastic syndrome or myeloproliferative disease associated with platelet-derived growth factor receptor gene re-arrangements, or G.) Aggressive systemic mastocytosis without the D816V c-KIT mutation or with c-KIT mutational status unknown
Age Restrictions	None
Prescriber Restrictions	None
Coverage Duration	12 months
Other Criteria	None
Indications	All Medically-accepted Indications.
Off-Label Uses	N/A
Part B Prerequisite	No

Formulary ID: 24449 version 13 Last Updated: 06/20/2024

Effective: 07/01/2024



IMBRUVICA

Products Affected

- IMBRUVICA ORAL CAPSULE
- IMBRUVICA ORAL SUSPENSION
- IMBRUVICA ORAL TABLET 140 MG, 280 MG, 420 MG

PA Criteria	Criteria Details
Exclusion Criteria	None
Required Medical Information	Diagnosis of one of the following A.) Chronic lymphocytic leukemia (CLL)/Small lymphocytic lymphoma (SLL), B.) Chronic lymphocytic leukemia (CLL)/Small lymphocytic lymphoma (SLL) with 17p deletion, C.) Waldenstrom's macroglobulinemia (WM), or D.) Chronic graft vs host disease (cGVHD) after failure of at least one first-line corticosteroid therapy
Age Restrictions	None
Prescriber Restrictions	None
Coverage Duration	12 months
Other Criteria	None
Indications	All Medically-accepted Indications.
Off-Label Uses	N/A
Part B Prerequisite	No

Formulary ID: 24449 version 13

Last Updated: 06/20/2024 Effective: 07/01/2024



INBRIJA

Products Affected

• INBRIJA

PA Criteria	Criteria Details
Exclusion Criteria	Any of the following A.) Concurrent use with nonselective monoamine oxidase inhibitors (MAOIs) (e.g. phenelzine and tranylcypromine), B.) Recent use (within 2 weeks) with a nonselective MAOI
Required Medical Information	Must meet all of the following A.) Diagnosis of Parkinson's disease and used for intermittent treatment of off episodes, and B.) Concurrent therapy with carbidopa/levodopa
Age Restrictions	18 years of age and older
Prescriber Restrictions	None
Coverage Duration	12 months
Other Criteria	None
Indications	All Medically-accepted Indications.
Off-Label Uses	N/A
Part B Prerequisite	No

Formulary ID: 24449 version 13

Last Updated: 06/20/2024 Effective: 07/01/2024



INCRELEX

Products Affected

INCRELEX

PA Criteria	Criteria Details
Exclusion Criteria	Any of the following A.) Active or suspected malignancy, B.) Use for growth promotion in patients with closed epiphyses, or C.) Intravenous administration
Required Medical Information	Diagnosis of one of the following A.) Severe primary insulin-like growth factor-1 (IGF-1) deficiency and utilized for pediatric treatment of growth failure, or B.) Growth hormone (GH) gene deletion and patient has developed neutralizing antibodies to GH and utilized for pediatric treatment of growth failure
Age Restrictions	None
Prescriber Restrictions	None
Coverage Duration	12 months
Other Criteria	None
Indications	All Medically-accepted Indications.
Off-Label Uses	N/A
Part B Prerequisite	No

Formulary ID: 24449 version 13 Last Updated: 06/20/2024

Effective: 07/01/2024



INLYTA

Products Affected

• INLYTA

PA Criteria	Criteria Details
Exclusion Criteria	None
Required Medical Information	Diagnosis of one of the following A.) Advanced renal cell carcinoma and patient failed one or more systemic therapies for renal cell carcinoma (e.g., sunitinib-, bevacizumab-, temsirolimus-, or cytokine-containing regimens), or B.) Advanced renal cell carcinoma and used as first-line therapy in combination with avelumab or pembrolizumab
Age Restrictions	18 years of age and older
Prescriber Restrictions	None
Coverage Duration	12 months
Other Criteria	None
Indications	All Medically-accepted Indications.
Off-Label Uses	N/A
Part B Prerequisite	No

Formulary ID: 24449 version 13 Last Updated: 06/20/2024

Effective: 07/01/2024



INQOVI

Products Affected

• INQOVI

PA Criteria	Criteria Details
Exclusion Criteria	None
Required Medical Information	Diagnosis of myelodysplastic syndromes (MDS), including previously treated and untreated, de novo and secondary MDS with the following French-American-British subtypes (refractory anemia, refractory anemia with ringed sideroblasts, refractory anemia with excess blasts, and chronic myelomonocytic leukemia [CMML]) and intermediate-1, intermediate-2, and high-risk International Prognostic Scoring System groups
Age Restrictions	18 years of age and older
Prescriber Restrictions	Prescribed by or in consultation with an oncologist
Coverage Duration	12 months
Other Criteria	None
Indications	All Medically-accepted Indications.
Off-Label Uses	N/A
Part B Prerequisite	No

Formulary ID: 24449 version 13 Last Updated: 06/20/2024

Effective: 07/01/2024



INREBIC

Products Affected

• INREBIC

PA Criteria	Criteria Details
Exclusion Criteria	None
Required Medical Information	Diagnosis of intermediate-2 or high-risk primary or secondary (post-polycythemia vera or post-essential thrombocythemia) myelofibrosis (MF).
Age Restrictions	18 years of age and older
Prescriber Restrictions	Prescribed by or in consultation with an oncologist or hematologist
Coverage Duration	12 months
Other Criteria	None
Indications	All Medically-accepted Indications.
Off-Label Uses	N/A
Part B Prerequisite	No

Formulary ID: 24449 version 13

Last Updated: 06/20/2024 Effective: 07/01/2024



INTRAROSA

Products Affected

• INTRAROSA

PA Criteria	Criteria Details
Exclusion Criteria	Any of the following A.) Vaginal bleeding or dysfunctional uterine bleeding of an undetermined origin, or B.) Known or suspected estrogendependent neoplasia
Required Medical Information	Diagnosis of one of the following A.) moderate to severe dyspareunia due to menopause, or B.) atrophic vaginitis due to menopause
Age Restrictions	18 years of age and older
Prescriber Restrictions	None
Coverage Duration	Initial: 3 months, Renewal: 12 months
Other Criteria	None
Indications	All Medically-accepted Indications.
Off-Label Uses	N/A
Part B Prerequisite	No

Formulary ID: 24449 version 13 Last Updated: 06/20/2024

Effective: 07/01/2024



ISTURISA

Products Affected

• ISTURISA ORAL TABLET 1 MG, 5 MG

PA Criteria	Criteria Details
Exclusion Criteria	None
Required Medical Information	Diagnosis of Cushing's disease in patients for whom pituitary surgery is not an option or has not been curative
Age Restrictions	18 years of age and older
Prescriber Restrictions	Prescribed by or in consultation with an endocrinologist
Coverage Duration	12 months
Other Criteria	None
Indications	All Medically-accepted Indications.
Off-Label Uses	N/A
Part B Prerequisite	No

Formulary ID: 24449 version 13 Last Updated: 06/20/2024

Effective: 07/01/2024



ITRACONAZOLE

Products Affected

• itraconazole oral

PA Criteria	Criteria Details
Exclusion Criteria	Any of the following A.) Ventricular dysfunction (e.g., congestive heart failure (CHF) or history of CHF), B.) Concurrent therapy with a CYP3A4 substrate (e.g., methadone, lovastatin, simvastatin, etc.), C.) Concurrent use of CYP2D6 inhibitors (e.g., bupropion, fluoxetine, paroxetine, quinidine, terbinafine), D.) Renal or hepatic impairment and concomitant use of colchicine, fesoterodine, solifenacin, or telithromycin, E.) Pregnancy
Required Medical Information	Diagnosis of one of the following A.) Systemic fungal infection (e.g., aspergillosis, histoplasmosis, blastomycosis), or B.) Onychomycosis confirmed by one of the following: positive potassium hydroxide (KOH) preparation, fungal culture, or nail biopsy
Age Restrictions	None
Prescriber Restrictions	None
Coverage Duration	6 months
Other Criteria	None
Indications	All Medically-accepted Indications.
Off-Label Uses	N/A
Part B Prerequisite	No

Formulary ID: 24449 version 13 Last Updated: 06/20/2024

Effective: 07/01/2024



ITRACONAZOLE SOLN

Products Affected

• itraconazole oral

PA Criteria	Criteria Details
Exclusion Criteria	Any of the following A.) Ventricular dysfunction (e.g., congestive heart failure (CHF) or history of CHF), B.) Concurrent therapy with a CYP3A4 substrate (e.g., methadone, lovastatin, simvastatin, etc.), C.) Concurrent use of CYP2D6 inhibitors (e.g., bupropion, fluoxetine, paroxetine, quinidine, terbinafine), D.) Renal or hepatic impairment and concomitant use of colchicine, fesoterodine, solifenacin, or telithromycin, E.) Pregnancy
Required Medical Information	Diagnosis of candidiasis (esophageal or oropharyngeal) that is refractory to treatment with fluconazole
Age Restrictions	None
Prescriber Restrictions	None
Coverage Duration	6 months
Other Criteria	None
Indications	All Medically-accepted Indications.
Off-Label Uses	N/A
Part B Prerequisite	No

Formulary ID: 24449 version 13 Last Updated: 06/20/2024

Effective: 07/01/2024



IVERMECTIN

Products Affected

• ivermectin oral

PA Criteria	Criteria Details
Exclusion Criteria	Prevention or treatment of COVID-19
Required Medical Information	Diagnosis of one of the following: A.) Strongyloidiasis of the intestinal tract or B.) Onchocerciasis
Age Restrictions	None
Prescriber Restrictions	None
Coverage Duration	1 month
Other Criteria	None
Indications	All Medically-accepted Indications.
Off-Label Uses	N/A
Part B Prerequisite	No

Formulary ID: 24449 version 13

Last Updated: 06/20/2024 Effective: 07/01/2024



IWILFIN

Products Affected

• IWILFIN

PA Criteria	Criteria Details
Exclusion Criteria	None
Required Medical Information	Diagnosis of high-risk neuroblastoma to be used to reduce the risk of relapse in adult and pediatric patients who have demonstrated at least a partial response to prior multiagent, multimodality therapy including anti-GD2 immunotherapy
Age Restrictions	None
Prescriber Restrictions	Prescribed by or in consultation with an oncologist
Coverage Duration	12 months
Other Criteria	None
Indications	All Medically-accepted Indications.
Off-Label Uses	N/A
Part B Prerequisite	No

Formulary ID: 24449 version 13

Last Updated: 06/20/2024 Effective: 07/01/2024



JAKAFI

Products Affected

JAKAFI

PA Criteria	Criteria Details
Exclusion Criteria	None
Required Medical Information	Diagnosis of one of the following A.) Intermediate or high-risk myelofibrosis, including primary myelofibrosis, post-polycythemia vera myelofibrosis and post-essential thrombocythemia myelofibrosis, B.) Polycythemia vera AND patient has had an inadequate response to or is intolerant of hydroxyurea, C.) Acute graft versus host disease AND disease is refractory to steroid therapy, or D.) Chronic graft-versus-host disease (cGVHD) after failure of corticosteroid therapy (alone or in combination with a calcineurin inhibitor) and up to one additional line of systemic therapy
Age Restrictions	None
Prescriber Restrictions	None
Coverage Duration	6 months
Other Criteria	None
Indications	All Medically-accepted Indications.
Off-Label Uses	N/A
Part B Prerequisite	No

Formulary ID: 24449 version 13

Last Updated: 06/20/2024 Effective: 07/01/2024



JAYPIRCA

Products Affected

JAYPIRCA

PA Criteria	Criteria Details
Exclusion Criteria	None
Required Medical Information	Diagnosis of one of the following A.) relapsed or refractory mantle cell lymphoma (MCL) and is being used after at least two lines of systemic therapy, including a BTK inhibitor or B.) chronic lymphocytic leukemia or small lymphocytic lymphoma who have received at least 2 prior lines of therapy, including a Bruton tyrosine kinase inhibitor and a B-cell lymphoma 2 inhibitor.
Age Restrictions	18 years of age and older
Prescriber Restrictions	Prescribed by or in consultation with an oncologist
Coverage Duration	12 months
Other Criteria	None
Indications	All Medically-accepted Indications.
Off-Label Uses	N/A
Part B Prerequisite	No

Formulary ID: 24449 version 13

Last Updated: 06/20/2024 Effective: 07/01/2024



JOENJA

Products Affected

JOENJA

PA Criteria	Criteria Details
Exclusion Criteria	None
Required Medical Information	Diagnosis of activated phosphoinositide 3-kinase (PI3K) delta syndrome
Age Restrictions	12 years of age and older
Prescriber Restrictions	None
Coverage Duration	12 months
Other Criteria	None
Indications	All Medically-accepted Indications.
Off-Label Uses	N/A
Part B Prerequisite	No

Formulary ID: 24449 version 13

Last Updated: 06/20/2024 Effective: 07/01/2024



JUXTAPID

Products Affected

• JUXTAPID ORAL CAPSULE 10 MG, 20 MG, 30 MG, 5 MG

PA Criteria	Criteria Details
Exclusion Criteria	Any of the following A.) Moderate to severe liver impairment or active liver disease including unexplained persistent abnormal liver function tests, B.) Pregnancy, or C.) Concomitant use with strong or moderate CYP3A4 inhibitors
Required Medical Information	Diagnosis of HoFH as confirmed by one of the following A.) Genetic confirmation of 2 mutations in the LDL receptor, ApoB, PCSK9, or LDL receptor adaptor protein 1 (LDLRAP1 or ARH), or B.) Both of the following 1.) Either untreated LDL-C greater than 500 mg/dL or treated LDL-C greater than 300 mg/dL, and 2.) Either xanthoma before 10 years of age or evidence of heterozygous FH in both parents
Age Restrictions	18 years of age and older
Prescriber Restrictions	None
Coverage Duration	Initial: 6 months, Renewal: 12 months
Other Criteria	None
Indications	All Medically-accepted Indications.
Off-Label Uses	N/A
Part B Prerequisite	No

Formulary ID: 24449 version 13 Last Updated: 06/20/2024

Effective: 07/01/2024



KALYDECO

Products Affected

KALYDECO

PA Criteria	Criteria Details
Exclusion Criteria	None
Required Medical Information	Diagnosis of cystic fibrosis (CF) and the patient has 1 mutation in the cystic fibrosis transmembrane conductance regulator (CFTR) gene that is responsive to ivacaftor potentiation based on clinical and/or in vitro assay data
Age Restrictions	None
Prescriber Restrictions	Prescribed by or in consultation with a pulmonologist or prescribing practitioner is from a CF center accredited by the Cystic Fibrosis Foundation
Coverage Duration	12 months
Other Criteria	None
Indications	All Medically-accepted Indications.
Off-Label Uses	N/A
Part B Prerequisite	No

Formulary ID: 24449 version 13 Last Updated: 06/20/2024

Effective: 07/01/2024



KESIMPTA

Products Affected

KESIMPTA

PA Criteria	Criteria Details
Exclusion Criteria	Active Hepatitis B infection
Required Medical Information	Diagnosis of one of the following A.) Relapsing forms of multiple sclerosis (e.g., clinically isolated syndrome, relapsing-remitting MS, active secondary progressive disease, or progressive-relapsing MS), or B.) Patient has experienced a first clinical episode and has MRI features consistent with multiple sclerosis
Age Restrictions	18 years of age and older
Prescriber Restrictions	Prescribed by or in consultation with a neurologist
Coverage Duration	12 months
Other Criteria	None
Indications	All Medically-accepted Indications.
Off-Label Uses	N/A
Part B Prerequisite	No

Formulary ID: 24449 version 13 Last Updated: 06/20/2024

Effective: 07/01/2024



KINERET

Products Affected

• KINERET SUBCUTANEOUS SOLUTION PREFILLED SYRINGE

PA Criteria	Criteria Details
Exclusion Criteria	None
Required Medical Information	Diagnosis of one of the following A.) Moderate to severe active rheumatoid arthritis and patient has trial and failure, contraindication, or intolerance to two preferred products, (i.e. Humira, Enbrel), B.) Cryopyrin-associated periodic syndromes (i.e., neonatal-onset multisystem inflammatory disease), or C.) Deficiency of interleukin-1 receptor antagonist
Age Restrictions	None
Prescriber Restrictions	None
Coverage Duration	12 months
Other Criteria	Screening for latent tuberculosis infection is required prior to initiation of treatment
Indications	All Medically-accepted Indications.
Off-Label Uses	N/A
Part B Prerequisite	No

Formulary ID: 24449 version 13 Last Updated: 06/20/2024

Effective: 07/01/2024



KISQALI

Products Affected

- KISQALI (200 MG DOSE)
- KISQALI (400 MG DOSE)

• KISQALI (600 MG DOSE)

PA Criteria	Criteria Details
Exclusion Criteria	None
Required Medical Information	Diagnosis of advanced or metastatic hormone receptor (HR)-positive, human epidermal growth factor receptor 2 (HER2)-negative breast cancer and one of the following A.) The patient is a pre-or perimenopausal woman or man and the requested drug will be used in combination with an aromatase inhibitor as initial endocrine-based therapy, B.) The patient is a postmenopausal woman or man, the requested drug will be used in combination with an aromatase inhibitor as initial endocrine-based therapy, and the patient has experienced disease progression, an intolerable adverse event, or contraindication to Ibrance (palbociclib) or Verzenio (abemaciclib), C.) The patient is a pre-or perimenopausal woman or man and the requested drug is being used with fulvestrant as initial endocrine-based therapy, or D.) The patient is a postmenopausal woman or man, the requested drug is being used following disease progression on endocrine therapy, and the patient has experienced disease progression, an intolerable adverse event, or contraindication to Ibrance (palbociclib) or Verzenio (abemaciclib)
Age Restrictions	18 years of age and older
Prescriber Restrictions	Prescribed by or in consultation with an oncologist
Coverage Duration	12 months
Other Criteria	None
Indications	All Medically-accepted Indications.
Off-Label Uses	N/A

Formulary ID: 24449 version 13 Last Updated: 06/20/2024

Effective: 07/01/2024



PA Criteria	Criteria Details
Part B Prerequisite	No

Formulary ID: 24449 version 13 Last Updated: 06/20/2024

Effective: 07/01/2024



KISQALI FEMARA

Products Affected

- KISQALI FEMARA (200 MG DOSE)
- KISQALI FEMARA (600 MG DOSE)
- KISQALI FEMARA (400 MG DOSE)

PA Criteria	Criteria Details
Exclusion Criteria	None
Required Medical Information	Diagnosis of advanced or metastatic hormone receptor (HR)-positive, human epidermal growth factor receptor 2 (HER2)-negative breast cancer and one of the following A.) The patient is pre-or perimenopausal woman or male and the requested drug will be used as initial endocrine-based therapy, B.) The patient is postmenopausal, the requested drug will be used as initial endocrine-based therapy, and the patient has experienced disease progression, an intolerable adverse event, or contraindication to Ibrance (palbociclib) or Verzenio (abemaciclib)
Age Restrictions	18 years of age and older
Prescriber Restrictions	Prescribed by or in consultation with an oncologist
Coverage Duration	12 months
Other Criteria	None
Indications	All Medically-accepted Indications.
Off-Label Uses	N/A
Part B Prerequisite	No

Formulary ID: 24449 version 13 Last Updated: 06/20/2024

Effective: 07/01/2024



KOSELUGO

Products Affected

KOSELUGO

PA Criteria	Criteria Details
Exclusion Criteria	None
Required Medical Information	Diagnosis of neurofibromatosis type 1 (NF1) in a patient who has symptomatic, inoperable plexiform neurofibromas (PN)
Age Restrictions	2 years of age to 17 years of age
Prescriber Restrictions	Prescribed by or in consultation with an oncologist
Coverage Duration	12 months
Other Criteria	None
Indications	All Medically-accepted Indications.
Off-Label Uses	N/A
Part B Prerequisite	No

Formulary ID: 24449 version 13

Last Updated: 06/20/2024 Effective: 07/01/2024



KRAZATI

Products Affected

KRAZATI

PA Criteria	Criteria Details
Exclusion Criteria	None
Required Medical Information	Diagnosis of KRAS G12C-mutated locally advanced or metastatic non- small cell lung cancer (NSCLC) as determined by an FDA-approved test and patient has received at least one prior systemic therapy
Age Restrictions	18 years of age and older
Prescriber Restrictions	Prescribed by or in consultation with an oncologist
Coverage Duration	12 months
Other Criteria	None
Indications	All Medically-accepted Indications.
Off-Label Uses	N/A
Part B Prerequisite	No

Formulary ID: 24449 version 13

Last Updated: 06/20/2024 Effective: 07/01/2024



LAPATINIB

Products Affected

• lapatinib ditosylate

PA Criteria	Criteria Details
Exclusion Criteria	None
Required Medical Information	Diagnosis of advanced or metastatic breast cancer with tumors that overexpress human epidermal growth factor receptor 2 (HER2) AND meets one of the following A.) Used in combination with capecitabine in a patient who has received prior therapy including an anthracycline, a taxane, and trastuzumab, OR B.) Used in combination with letrozole in a postmenopausal female for whom hormonal therapy is indicated
Age Restrictions	18 years of age and older
Prescriber Restrictions	Prescribed by or in consultation with an oncologist
Coverage Duration	12 months
Other Criteria	None
Indications	All Medically-accepted Indications.
Off-Label Uses	N/A
Part B Prerequisite	No

Formulary ID: 24449 version 13 Last Updated: 06/20/2024

Effective: 07/01/2024



LENALIDOMIDE

Products Affected

• lenalidomide

PA Criteria	Criteria Details
ra Criteria	Criteria Details
Exclusion Criteria	Pregnancy
Required Medical Information	Diagnosis of one of the following A.) Multiple myeloma and medication will be used in combination with dexamethasone, B.) Autologous hematopoietic stem-cell transplantation (HSCT) in multiple myeloma patients, C.) Transfusion-dependent anemia due to low- or intermediate-1-risk myelodysplastic syndrome (MDS) associated with a deletion 5q cytogenetic abnormality or without additional cytogenetic abnormalities, D.) Mantle cell lymphoma whose disease has relapsed or progressed after two prior therapies, one of which included bortezomib, E.) Follicular lymphoma and used in combination with rituximab, or F.) Marginal zone lymphoma and used in combination with rituximab
Age Restrictions	None
Prescriber Restrictions	None
Coverage Duration	12 months
Other Criteria	None
Indications	All Medically-accepted Indications.
Off-Label Uses	N/A
Part B Prerequisite	No

Formulary ID: 24449 version 13 Last Updated: 06/20/2024

Effective: 07/01/2024



LENVIMA

Products Affected

- LENVIMA (10 MG DAILY DOSE)
- LENVIMA (12 MG DAILY DOSE)
- LENVIMA (14 MG DAILY DOSE)
- LENVIMA (18 MG DAILY DOSE)
- LENVIMA (20 MG DAILY DOSE)
- LENVIMA (24 MG DAILY DOSE)
- LENVIMA (4 MG DAILY DOSE)
- LENVIMA (8 MG DAILY DOSE)

PA Criteria	Criteria Details
Exclusion Criteria	None
Required Medical Information	Diagnosis of one of the following A.) Locally recurrent or metastatic, progressive, radioactive iodine-refractory differentiated thyroid cancer, B.) Advanced renal cell carcinoma, in combination with everolimus, following one prior anti-angiogenic therapy, C.) Unresectable hepatocellular carcinoma, first-line therapy, D.) Advanced endometrial carcinoma that is not microsatellite instability-high or mismatch repair deficient, in combination with pembrolizumab, when disease has progressed following prior systemic therapy and patient is not a candidate for curative surgery or radiation, or E.) Advanced renal cell carcinoma, in combination with pembrolizumab and used as first-line therapy
Age Restrictions	18 years of age and older
Prescriber Restrictions	None
Coverage Duration	12 months
Other Criteria	None
Indications	All Medically-accepted Indications.
Off-Label Uses	N/A
Part B Prerequisite	No

Formulary ID: 24449 version 13 Last Updated: 06/20/2024

Effective: 07/01/2024



LEUKINE

Products Affected

• LEUKINE INJECTION SOLUTION RECONSTITUTED

PA Criteria	Criteria Details
Exclusion Criteria	None
Required Medical Information	Diagnosis of one of the following A.) Patient has undergone allogeneic or autologous bone marrow transplant (BMT) and engraftment is delayed or failed, B.) Patient is undergoing autologous peripheral-blood progenitor cell transplant to mobilize progenitor cells for collection by leukapheresis, C.) Medication will be used for myeloid reconstitution after an autologous or allogeneic BMT, D.) Patient has acute myeloid leukemia and administration will be after completion of induction chemotherapy, E.) Hematopoietic subsyndrome of acute radiation syndrome (H-ARS) or F.) Autologous peripheral blood stem cell transplant, Following myeloablative chemotherapy.
Age Restrictions	None
Prescriber Restrictions	None
Coverage Duration	12 months
Other Criteria	None
Indications	All Medically-accepted Indications.
Off-Label Uses	N/A
Part B Prerequisite	No

Formulary ID: 24449 version 13 Last Updated: 06/20/2024

Effective: 07/01/2024



LEUPROLIDE

Products Affected

- ELIGARD
- *leuprolide acetate (3 month)*
- leuprolide acetate injection
- LUPRON DEPOT (1-MONTH)
- LUPRON DEPOT (3-MONTH)
- LUPRON DEPOT (4-MONTH)

- LUPRON DEPOT (6-MONTH)
- LUPRON DEPOT-PED (1-MONTH) INTRAMUSCULAR KIT 7.5 MG
- LUPRON DEPOT-PED (3-MONTH) INTRAMUSCULAR KIT 11.25 MG
- LUPRON DEPOT-PED (6-MONTH)

PA Criteria	Criteria Details
Exclusion Criteria	Any of the following A.) Pregnancy, B.) Undiagnosed abnormal uterine bleeding
Required Medical Information	Must meet one of the following: 1.) Eligard only: Advanced or metastatic prostate cancer, 2.) For Lupron depot and leuprolide products only: A.) Advanced or metastatic prostate cancer and patient has failed or is intolerant to Eligard (7.5 mg 1-month, 22.5 mg 3-month, 30 mg 4-month, & 45 mg 6-month depots only), B.) Endometriosis (3.75 mg 1-month & 11.25 mg 3-month depots only), C.) Anemia due to uterine leiomyomata (Fibroids) (3.75 mg 1-month &11.25 mg 3-month depots only) and patient is preoperative, or D.) Central precocious puberty (idiopathic or neurogenic) in children
Age Restrictions	None
Prescriber Restrictions	None
Coverage Duration	12 months
Other Criteria	B vs D determination required per CMS guidance
Indications	All Medically-accepted Indications.
Off-Label Uses	N/A
Part B Prerequisite	No

Formulary ID: 24449 version 13 Last Updated: 06/20/2024

Effective: 07/01/2024



LIDOCAINE PATCH

Products Affected

• lidocaine external patch 5 %

PA Criteria	Criteria Details
Exclusion Criteria	None
Required Medical Information	Diagnosis of one of the following A.) Pain associated with diabetic neuropathy, B.) Pain associated with cancer-related neuropathy, C.) Post-herpetic neuralgia, D.) Chronic back pain, or E.) Osteoarthritis of the knee or hip
Age Restrictions	None
Prescriber Restrictions	None
Coverage Duration	12 months
Other Criteria	None
Indications	All Medically-accepted Indications.
Off-Label Uses	N/A
Part B Prerequisite	No

Formulary ID: 24449 version 13

Last Updated: 06/20/2024 Effective: 07/01/2024



LINEZOLID

Products Affected

• linezolid intravenous solution 600 mg/300ml

• linezolid oral tablet

ing, soone	
PA Criteria	Criteria Details
Exclusion Criteria	Any of the following A.) Concomitant use of an MAOI, or B.) Use within 14 days of discontinuing an MAOI
Required Medical Information	Diagnosis of one of the following A.) Community acquired pneumonia, B.) Hospital-acquired pneumonia, C.) Vancomycin-resistant Enterococcus faecium infection, D.) Complicated skin and skin structure infections, or E.) Uncomplicated skin and skin structure infections
Age Restrictions	None
Prescriber Restrictions	None
Coverage Duration	1 month
Other Criteria	IV formulation: B vs D determination required per CMS guidance
Indications	All Medically-accepted Indications.
Off-Label Uses	N/A
Part B Prerequisite	No

Formulary ID: 24449 version 13 Last Updated: 06/20/2024

Effective: 07/01/2024



LIVMARLI

Products Affected

LIVMARLI

PA Criteria	Criteria Details
ra Cilicila	Criteria Details
Exclusion Criteria	None
Required Medical Information	Diagnosis of one of the following A.) Cholestatic pruritus in patients with Alagille syndrome (ALGS), or B.) Cholestatic pruritus in patients 5 years and older with progressive familial intrahepatic cholestasis (PFIC), but patient is not in a subgroup of PFIC type 2 patients with specific ABCB11 variants resulting in nonfunctional or complete absence of bile salt export pump protein
Age Restrictions	None
Prescriber Restrictions	None
Coverage Duration	12 months
Other Criteria	None
Indications	All Medically-accepted Indications.
Off-Label Uses	N/A
Part B Prerequisite	No

Formulary ID: 24449 version 13

Last Updated: 06/20/2024 Effective: 07/01/2024



LIVTENCITY

Products Affected

• LIVTENCITY

PA Criteria	Criteria Details
Exclusion Criteria	None
Required Medical Information	Diagnosis of post-transplant cytomegalovirus (CMV) infection/disease that is refractory to treatment (with or without genotypic resistance) with ganciclovir, valganciclovir, cidofovir or foscarnet
Age Restrictions	12 years of age and older
Prescriber Restrictions	None
Coverage Duration	12 months
Other Criteria	None
Indications	All Medically-accepted Indications.
Off-Label Uses	N/A
Part B Prerequisite	No

Formulary ID: 24449 version 13

Last Updated: 06/20/2024 Effective: 07/01/2024



LONSURF

Products Affected

• LONSURF

PA Criteria	Criteria Details
Exclusion Criteria	None
Required Medical Information	Diagnosis of one of the following A.) Metastatic colorectal cancer, previously treated with fluoropyrimidine, oxaliplatin, and irinotecan-based regimens, an anti-VEGF therapy, and if RAS wild-type, an anti-EGFR therapy, or B.) Metastatic gastric or gastroesophageal junction adenocarcinoma previously treated with at least 2 prior lines of chemotherapy that included a fluoropyrimidine, a platinum, either a taxane or irinotecan and if appropriate, HER2/neu-targeted therapy
Age Restrictions	18 years of age and older
Prescriber Restrictions	None
Coverage Duration	12 months
Other Criteria	None
Indications	All Medically-accepted Indications.
Off-Label Uses	N/A
Part B Prerequisite	No

Formulary ID: 24449 version 13

Last Updated: 06/20/2024 Effective: 07/01/2024



LORBRENA

Products Affected

LORBRENA

PA Criteria	Criteria Details
Exclusion Criteria	Concomitant use with strong CYP3A4 inducers
Required Medical Information	Diagnosis of anaplastic lymphoma kinase (ALK)-positive metastatic non-small cell lung cancer (NSCLC) as detected by an FDA-approved test
Age Restrictions	18 years of age and older
Prescriber Restrictions	Prescribed by or in consultation with an oncologist
Coverage Duration	12 months
Other Criteria	None
Indications	All Medically-accepted Indications.
Off-Label Uses	N/A
Part B Prerequisite	No

Formulary ID: 24449 version 13

Last Updated: 06/20/2024 Effective: 07/01/2024



LUMAKRAS

Products Affected

• LUMAKRAS

PA Criteria	Criteria Details
Exclusion Criteria	None
Required Medical Information	Diagnosis of KRAS G12C-mutated locally advanced or metastatic non-small cell lung cancer (NSCLC) as determined by an FDA-approved test, and patient has received at least one prior systemic therapy (e.g., immune checkpoint inhibitor, platinum-based chemotherapy)
Age Restrictions	18 years of age and older
Prescriber Restrictions	Prescribed by or in consultation with an oncologist
Coverage Duration	12 months
Other Criteria	None
Indications	All Medically-accepted Indications.
Off-Label Uses	N/A
Part B Prerequisite	No

Formulary ID: 24449 version 13

Last Updated: 06/20/2024 Effective: 07/01/2024



LUPKYNIS

Products Affected

LUPKYNIS

PA Criteria	Criteria Details
Exclusion Criteria	Concomitant use of strong CYP3A4 inhibitors (e.g., ketoconazole, itraconazole, clarithromycin)
Required Medical Information	Initial: Diagnosis of systemic lupus erythematosus (SLE) with active lupus nephritis (LN) Classes III, IV, V (alone or in combination), and all of the following: 1.) Baseline renal function of 45 mL/min/1.73 m2 or greater, 2.) Will be used in combination with a background immunosuppressive therapy regimen (e.g. mycophenolate, oral steroids, etc). Renewal: Documentation of positive clinical response to therapy
Age Restrictions	18 years of age and older
Prescriber Restrictions	Prescribed by or in consultation with a rheumatologist or nephrologist
Coverage Duration	Initial: 12 months, Renewal: 12 months
Other Criteria	None
Indications	All Medically-accepted Indications.
Off-Label Uses	N/A
Part B Prerequisite	No

Formulary ID: 24449 version 13 Last Updated: 06/20/2024

Effective: 07/01/2024



LYNPARZA

Products Affected

• LYNPARZA ORAL TABLET

PA Criteria	Criteria Details
Exclusion Criteria	None
Required Medical Information	Diagnosis of one of the following A.) HER2-negative, deleterious or suspected deleterious germline BRCA mutated high-risk early or metastatic breast cancer AND patient has been previously treated with chemotherapy in neoadjuvant, adjuvant, or metastatic setting, B.) Recurrent epithelial ovarian cancer, recurrent fallopian tube cancer, or recurrent primary peritoneal cancer AND used for maintenance treatment in patients who are in complete or partial response to platinum-based chemotherapy (e.g. cisplatin, carboplatin), C.) Deleterious or suspected deleterious germline or somatic BRCA-mutated (gBRCAm or sBRCAm) epithelial ovarian, fallopian tube, or primary peritoneal cancer in patients with complete or partial response to first-line platinum-based chemotherapy, D.) Deleterious or suspected deleterious germline BRCA-mutated metastatic pancreatic adenocarcinoma and disease has not progressed on at least 16 weeks of a first-line platinum-based chemotherapy regimen, E.) Advanced epithelial ovarian, fallopian tube, or primary peritoneal cancer in patients who are in complete or partial response to first-line platinum-based chemotherapy and whose cancer is associated with homologous recombination deficiency positive status defined by either a deleterious or suspected deleterious BRCA-mutation and/or genomic instability AND are using in combination with bevacizumab for maintenance treatment, F.) Deleterious or suspected deleterious germline or somatic homologous recombination repair gene mutated metastatic castration-resistant prostate cancer in patients who have progressed following prior treatment with enzalutamide or abiraterone, or G.) Deleterious or suspected deleterious BRCA-mutated metastatic castration-resistant prostate cancer in combination with abiraterone and prednisone or prednisolone
Age Restrictions	18 years of age and older
Prescriber Restrictions	Prescribed by or in consultation with an oncologist

Formulary ID: 24449 version 13 Last Updated: 06/20/2024

Effective: 07/01/2024



PA Criteria	Criteria Details
Coverage Duration	12 months
Other Criteria	None
Indications	All Medically-accepted Indications.
Off-Label Uses	N/A
Part B Prerequisite	No

Formulary ID: 24449 version 13 Last Updated: 06/20/2024

Effective: 07/01/2024



LYTGOBI

Products Affected

- LYTGOBI (12 MG DAILY DOSE)
- LYTGOBI (20 MG DAILY DOSE)
- LYTGOBI (16 MG DAILY DOSE)

PA Criteria	Criteria Details
Exclusion Criteria	None
Required Medical Information	Diagnosis of unresectable, locally advanced or metastatic intrahepatic cholangiocarcinoma harboring fibroblast growth factor receptor 2 (FGFR2) gene fusions or other rearrangements and previously treated
Age Restrictions	18 years of age and older
Prescriber Restrictions	Prescribed by or in consultation with an oncologist
Coverage Duration	12 months
Other Criteria	None
Indications	All Medically-accepted Indications.
Off-Label Uses	N/A
Part B Prerequisite	No

Formulary ID: 24449 version 13

Last Updated: 06/20/2024 Effective: 07/01/2024

 $H9306_24_DRS_09\text{-}001_OE_C$



MATULANE

Products Affected

MATULANE

PA Criteria	Criteria Details
Exclusion Criteria	Inadequate marrow reserve
Required Medical Information	Diagnosis of Hodgkin's Disease, Stages III and IV and used in combination with other anticancer drugs
Age Restrictions	None
Prescriber Restrictions	Prescribed by or in consultation with an oncologist
Coverage Duration	12 months
Other Criteria	None
Indications	All Medically-accepted Indications.
Off-Label Uses	N/A
Part B Prerequisite	No

Formulary ID: 24449 version 13 Last Updated: 06/20/2024

Effective: 07/01/2024



MAYZENT

Products Affected

MAYZENT

• MAYZENT STARTER PACK

PA Criteria	Criteria Details
Exclusion Criteria	Any of the following A.) CYP2C9*3/*3 genotype, B.) In the last 6 months experienced myocardial infarction, unstable angina, stroke, TIA, decompensated heart failure requiring hospitalization, Class III-IV heart failure, or C.) Presence of Mobitz type II second-degree, third-degree AV block, or sick sinus syndrome, unless patient has a functioning pacemaker
Required Medical Information	Diagnosis of relapsing forms of multiple sclerosis, including clinically isolated syndrome, relapsing-remitting disease, or active secondary progressive disease and the following A.) Patients with relapsing forms of multiple sclerosis have history of/or contraindication to Avonex, Betaseron, Copaxone/Glatiramer, Gilenya/Fingolimod, or Dimethyl Fumarate
Age Restrictions	18 years of age and older
Prescriber Restrictions	Prescribed by or in consultation with a neurologist
Coverage Duration	12 months
Other Criteria	None
Indications	All Medically-accepted Indications.
Off-Label Uses	N/A
Part B Prerequisite	No

Formulary ID: 24449 version 13 Last Updated: 06/20/2024

Effective: 07/01/2024



MEKINIST

Products Affected

MEKINIST

PA Criteria	Criteria Details
Exclusion Criteria	None
Required Medical Information	Diagnosis of one of the following A.) Locally advanced or metastatic anaplastic thyroid cancer (ATC) with BRAF V600E mutation and used in combination with dabrafenib and no locoregional treatment options, B.) Malignant melanoma with lymph node involvement and following complete resection with BRAF V600E or V600K mutations and used in combination with dabrafenib, C.) Unresectable or metastatic malignant melanoma with BRAF V600E or V600K mutations and used in combination with dabrafenib or as monotherapy, D.) Metastatic non-small cell lung cancer, with BRAF V600E mutation, in combination with dabrafenib, E.) Unresectable or metastatic solid tumors with BRAF V600E mutation, in combination with dabrafenib, and have progressed following prior treatment and have no satisfactory alternative treatment options, F.) Low-grade glioma with a BRAF V600E mutation and require systemic therapy, in combination with dabrafenib
Age Restrictions	None
Prescriber Restrictions	Prescribed by or in consultation with an oncologist
Coverage Duration	12 months
Other Criteria	None
Indications	All Medically-accepted Indications.
Off-Label Uses	N/A
Part B Prerequisite	No

Formulary ID: 24449 version 13 Last Updated: 06/20/2024

Effective: 07/01/2024



MEKTOVI

Products Affected

MEKTOVI

PA Criteria	Criteria Details
Exclusion Criteria	None
Required Medical Information	Diagnosis of one of the following A.) Unresectable or metastatic malignant melanoma with documented BRAF V600E or V600K mutation as detected by an FDA approved test AND used in combination with encorafenib or B.) Metastatic non-small cell lung cancer with a BRAF V600E mutation as detected by an FDA-approved test AND used in combination with encorafenib
Age Restrictions	18 years of age and older
Prescriber Restrictions	Prescribed by or in consultation with an oncologist
Coverage Duration	12 months
Other Criteria	None
Indications	All Medically-accepted Indications.
Off-Label Uses	N/A
Part B Prerequisite	No

Formulary ID: 24449 version 13 Last Updated: 06/20/2024

Effective: 07/01/2024



MIFEPRISTONE

Products Affected

• mifepristone oral tablet 300 mg

PA Criteria	Criteria Details
Exclusion Criteria	Any of the following A.) pregnancy, B.) coadministration with simvastatin, lovastatin, or CYP3A substrates with narrow therapeutic ranges, C.) concomitant treatment with systemic corticosteroids for serious medical conditions or illnesses, D.) history of unexplained vaginal bleeding, E.) endometrial hyperplasia with atypia or endometrial carcinoma
Required Medical Information	Diagnosis of endogenous Cushing syndrome in patients with type 2 diabetes mellitus or glucose intolerance and must meet all of the following 1.) Used to control hyperglycemia secondary to hypercortisolism, and 2.) Patient has failed or is not a candidate for surgery
Age Restrictions	18 years of age and older
Prescriber Restrictions	Prescribed by or in consultation with an endocrinologist
Coverage Duration	12 months
Other Criteria	None
Indications	All Medically-accepted Indications.
Off-Label Uses	N/A
Part B Prerequisite	No

Formulary ID: 24449 version 13 Last Updated: 06/20/2024

Effective: 07/01/2024



MIGLUSTAT

Products Affected

• miglustat

• YARGESA

PA Criteria	Criteria Details
Exclusion Criteria	None
Required Medical Information	Diagnosis of mild to moderate type 1 Gaucher disease and patient is not a candidate for enzyme replacement therapy
Age Restrictions	18 years of age and older
Prescriber Restrictions	None
Coverage Duration	12 months
Other Criteria	None
Indications	All Medically-accepted Indications.
Off-Label Uses	N/A
Part B Prerequisite	No

Formulary ID: 24449 version 13 Last Updated: 06/20/2024

Effective: 07/01/2024



MS INTERFERONS

Products Affected

- AVONEX PEN INTRAMUSCULAR AUTO-INJECTOR KIT
- AVONEX PREFILLED INTRAMUSCULAR PREFILLED SYRINGE KIT
- BETASERON SUBCUTANEOUS KIT

PA Criteria	Criteria Details
Exclusion Criteria	None
Required Medical Information	Diagnosis of one of the following A.) Relapsing forms of multiple sclerosis (e.g., clinically isolated syndrome, relapsing-remitting MS, active secondary progressive disease, or progressive-relapsing MS), or B.) Patient has experienced a first clinical episode and has MRI features consistent with multiple sclerosis
Age Restrictions	18 years of age and older
Prescriber Restrictions	Prescribed by or in consultation with a neurologist
Coverage Duration	12 months
Other Criteria	None
Indications	All Medically-accepted Indications.
Off-Label Uses	N/A
Part B Prerequisite	No

Formulary ID: 24449 version 13 Last Updated: 06/20/2024

Effective: 07/01/2024



NAMZARIC

Products Affected

NAMZARIC

PA Criteria	Criteria Details
Exclusion Criteria	Any of the following A.) Hypersensitivity to memantine, donepezil, or excipients, or B.) Hypersensitivity to piperidine derivatives
Required Medical Information	Diagnosis of moderate to severe dementia associated with Alzheimer's disease
Age Restrictions	18 years of age and older
Prescriber Restrictions	None
Coverage Duration	12 months
Other Criteria	None
Indications	All Medically-accepted Indications.
Off-Label Uses	N/A
Part B Prerequisite	No

Formulary ID: 24449 version 13 Last Updated: 06/20/2024

Effective: 07/01/2024



NERLYNX

Products Affected

NERLYNX

PA Criteria	Criteria Details
Exclusion Criteria	None
Required Medical Information	Diagnosis of one of the following A.) Early stage HER2-positive breast cancer and used following adjuvant trastuzumab therapy, or B.) Advanced or metastatic HER2-positive breast cancer, used in combination with capecitabine, AND patient has received 2 or more prior anti-HER2-based regimens in the metastatic setting
Age Restrictions	18 years of age and older
Prescriber Restrictions	Prescribed by or in consultation with an oncologist
Coverage Duration	12 months
Other Criteria	None
Indications	All Medically-accepted Indications.
Off-Label Uses	N/A
Part B Prerequisite	No

Formulary ID: 24449 version 13 Last Updated: 06/20/2024

Effective: 07/01/2024



NINLARO

Products Affected

NINLARO

PA Criteria	Criteria Details
Exclusion Criteria	None
Required Medical Information	Diagnosis of multiple myeloma, used in combination with lenalidomide and dexamethasone, AND patient has history of at least 1 prior therapy
Age Restrictions	18 years of age and older
Prescriber Restrictions	None
Coverage Duration	12 months
Other Criteria	None
Indications	All Medically-accepted Indications.
Off-Label Uses	N/A
Part B Prerequisite	No

Formulary ID: 24449 version 13 Last Updated: 06/20/2024

Effective: 07/01/2024



NITISINONE

Products Affected

• nitisinone

PA Criteria	Criteria Details
Exclusion Criteria	None
Required Medical Information	Diagnosis of hereditary tyrosinemia type 1 confirmed by one of the following A.) Biochemical testing (e.g., detection of succinylacetone in urine), or B.) DNA testing (mutation analysis)
Age Restrictions	None
Prescriber Restrictions	None
Coverage Duration	12 months
Other Criteria	None
Indications	All Medically-accepted Indications.
Off-Label Uses	N/A
Part B Prerequisite	No

Formulary ID: 24449 version 13

Last Updated: 06/20/2024 Effective: 07/01/2024



NUBEQA

Products Affected

NUBEQA

PA Criteria	Criteria Details
Exclusion Criteria	None
Required Medical Information	Diagnosis of one of the following A.) Non-metastatic, castration-resistant prostate cancer (nmCRPC) or B.) Metastatic hormone-sensitive prostate cancer in combination with docetaxel. For treatment of nmCRPC, one of the following applies 1.) Used in combination with a gonadotropin-releasing hormone (GnRH) analog or 2) Patient has received bilateral orchiectomy
Age Restrictions	18 years of age and older
Prescriber Restrictions	Prescribed by or in consultation with an oncologist or urologist
Coverage Duration	12 months
Other Criteria	None
Indications	All Medically-accepted Indications.
Off-Label Uses	N/A
Part B Prerequisite	No

Formulary ID: 24449 version 13

Last Updated: 06/20/2024 Effective: 07/01/2024



NUCALA

Products Affected

• NUCALA

PA Criteria	Criteria Details
Exclusion Criteria	None
Required Medical Information	Diagnosis of one of the following A.) Severe asthma with eosinophilic phenotype and used as an adjunct treatment, B.) Eosinophilic granulomatosis with polyangiitis (EGPA), C.) Hypereosinophilic syndrome lasting at least 6 months without an identifiable non-hematologic secondary cause, or D.) Chronic rhinosinusitis with nasal polyps and used as an adjunct treatment
Age Restrictions	6 years of age and older
Prescriber Restrictions	None
Coverage Duration	12 months
Other Criteria	None
Indications	All Medically-accepted Indications.
Off-Label Uses	N/A
Part B Prerequisite	No

Formulary ID: 24449 version 13 Last Updated: 06/20/2024

Effective: 07/01/2024



NUEDEXTA

Products Affected

NUEDEXTA

PA Criteria	Criteria Details
Exclusion Criteria	Any of the following A.) History of prolonged QT interval, congenital long QT syndrome or Torsades de pointes, B.) Heart failure, C.) Complete AV block without an implanted pacemaker or high risk of complete AV block, D.) Concomitant use with quinidine, quinine, mefloquine, or drugs that prolong QT interval and are metabolized by CYP2D6 (e.g., thioridazine, pimozide), E.) Concomitant use with MAOIs or within 14 days of MAOI therapy, F.) History of quinine-, mefloquine-, or quinidine-induced thrombocytopenia, bone marrow depression, or lupus-like syndrome
Required Medical Information	Diagnosis of pseudobulbar affect
Age Restrictions	18 years of age and older
Prescriber Restrictions	None
Coverage Duration	12 months
Other Criteria	None
Indications	All Medically-accepted Indications.
Off-Label Uses	N/A
Part B Prerequisite	No

Formulary ID: 24449 version 13 Last Updated: 06/20/2024

Effective: 07/01/2024



NUPLAZID

Products Affected

• NUPLAZID ORAL CAPSULE

NUPLAZID ORAL TABLET 10 MG

PA Criteria	Criteria Details
Exclusion Criteria	None
Required Medical Information	Diagnosis of Parkinson's disease and both of the following apply A.) Used for treatment of hallucinations and/or delusions associated with Parkinson's disease psychosis, and B.) Diagnosis of Parkinson's disease was made prior to the onset of psychotic symptoms
Age Restrictions	18 years of age and older
Prescriber Restrictions	None
Coverage Duration	12 months
Other Criteria	None
Indications	All Medically-accepted Indications.
Off-Label Uses	N/A
Part B Prerequisite	No

Formulary ID: 24449 version 13

Last Updated: 06/20/2024 Effective: 07/01/2024



OCTREOTIDE

Products Affected

• octreotide acetate injection solution 100 mcg/ml, 1000 mcg/ml, 200 mcg/ml, 50 mcg/ml, 500 mcg/ml

PA Criteria	Criteria Details
Exclusion Criteria	None
Required Medical Information	Diagnosis of one of the following A.) Acromegaly confirmed by high pretreatment insulin-like growth factor-1 (IGF-1) level for age and/or gender based on the laboratory reference range and patient has had inadequate response to or is ineligible for surgery, radiation, or bromocriptine mesylate, or B.) Metastatic carcinoid syndrome with associated diarrhea or flushing, or C.) Vasoactive intestinal peptidesecreting tumors (VIPomas) with associated diarrhea
Age Restrictions	18 years of age and older
Prescriber Restrictions	None
Coverage Duration	12 months
Other Criteria	B vs D determination required per CMS guidance
Indications	All Medically-accepted Indications.
Off-Label Uses	N/A
Part B Prerequisite	No

Formulary ID: 24449 version 13 Last Updated: 06/20/2024

Effective: 07/01/2024



ODOMZO

Products Affected

ODOMZO

PA Criteria	Criteria Details
Exclusion Criteria	Pregnancy
Required Medical Information	Diagnosis of locally advanced basal cell carcinoma of the skin and one of the following A.) Cancer has recurred following surgery or radiation therapy, B.) Patient is not a candidate for surgery or radiation therapy.
Age Restrictions	18 years of age and older
Prescriber Restrictions	None
Coverage Duration	12 months
Other Criteria	None
Indications	All Medically-accepted Indications.
Off-Label Uses	N/A
Part B Prerequisite	No

Formulary ID: 24449 version 13 Last Updated: 06/20/2024

Effective: 07/01/2024



OFEV

Products Affected

OFEV

PA Criteria	Criteria Details
Exclusion Criteria	None
Required Medical Information	Diagnosis of one of the following A.) Idiopathic pulmonary fibrosis (IPF), B.) Systemic sclerosis-associated interstitial lung disease (ILD), or C.) Chronic fibrosing interstitial lung disease with a progressive phenotype
Age Restrictions	18 years of age and older
Prescriber Restrictions	Prescribed by or in consultation with a pulmonologist
Coverage Duration	12 months
Other Criteria	None
Indications	All Medically-accepted Indications.
Off-Label Uses	N/A
Part B Prerequisite	No

Formulary ID: 24449 version 13

Last Updated: 06/20/2024 Effective: 07/01/2024



OGSIVEO

Products Affected

• OGSIVEO ORAL TABLET 50 MG

PA Criteria	Criteria Details
Exclusion Criteria	None
Required Medical Information	Diagnosis of progressing desmoid tumors who require systemic treatment
Age Restrictions	18 years or older
Prescriber Restrictions	Prescribed by or in consultation with an oncologist
Coverage Duration	12 months
Other Criteria	None
Indications	All Medically-accepted Indications.
Off-Label Uses	N/A
Part B Prerequisite	No

Formulary ID: 24449 version 13

Last Updated: 06/20/2024 Effective: 07/01/2024



OJJAARA

Products Affected

OJJAARA

PA Criteria	Criteria Details
Exclusion Criteria	None
Required Medical Information	Diagnosis of intermediate or high-risk myelofibrosis (MF), including primary MF or secondary MF [postpolycythemia vera (PV) and postessential thrombocythemia (ET)], in adults with anemia.
Age Restrictions	18 years of age and older
Prescriber Restrictions	Prescribed by or in consultation with an oncologist or hematologist
Coverage Duration	12 months
Other Criteria	None
Indications	All Medically-accepted Indications.
Off-Label Uses	N/A
Part B Prerequisite	No

Formulary ID: 24449 version 13

Last Updated: 06/20/2024 Effective: 07/01/2024



ONUREG

Products Affected

• ONUREG

PA Criteria	Criteria Details
Exclusion Criteria	None
Required Medical Information	Diagnosis of acute myeloid leukemia (AML) used in maintenance treatment for adult patients who achieved first complete remission (CR) or complete remission with incomplete blood count recovery (CRi) following intensive induction chemotherapy and are not able to complete intensive curative therapy
Age Restrictions	18 years of age and older
Prescriber Restrictions	Prescribed by or in consultation with an oncologist or hematologist
Coverage Duration	12 months
Other Criteria	None
Indications	All Medically-accepted Indications.
Off-Label Uses	N/A
Part B Prerequisite	No

Formulary ID: 24449 version 13 Last Updated: 06/20/2024

Effective: 07/01/2024



OPSUMIT

Products Affected

• OPSUMIT

PA Criteria	Criteria Details
Exclusion Criteria	Pregnancy
Required Medical Information	Diagnosis of pulmonary arterial hypertension (WHO Group I), confirmed by right heart catheterization or Doppler echocardiogram if patient is unable to undergo a right heart catheterization (e. g., patient is frail, elderly, etc.)
Age Restrictions	18 years of age and older
Prescriber Restrictions	Prescribed by or in consultation with a cardiologist or pulmonologist
Coverage Duration	Initial: 6 months, Renewal: 12 months
Other Criteria	None
Indications	All Medically-accepted Indications.
Off-Label Uses	N/A
Part B Prerequisite	No

Formulary ID: 24449 version 13 Last Updated: 06/20/2024

Effective: 07/01/2024



ORGOVYX

Products Affected

ORGOVYX

PA Criteria	Criteria Details
Exclusion Criteria	None
Required Medical Information	Diagnosis of advanced prostate cancer
Age Restrictions	18 years of age and older
Prescriber Restrictions	Prescribed by or in consultation with an oncologist or urologist
Coverage Duration	12 months
Other Criteria	None
Indications	All Medically-accepted Indications.
Off-Label Uses	N/A
Part B Prerequisite	No

Formulary ID: 24449 version 13 Last Updated: 06/20/2024

Effective: 07/01/2024



ORKAMBI

Products Affected

ORKAMBI

PA Criteria	Criteria Details
Exclusion Criteria	None
Required Medical Information	Diagnosis of cystic fibrosis (CF) with documented homozygous F508del mutation confirmed by FDA-approved CF mutation test
Age Restrictions	None
Prescriber Restrictions	Prescribed by or in consultation with a pulmonologist or prescribing practitioner is from a CF center accredited by the Cystic Fibrosis Foundation
Coverage Duration	12 months
Other Criteria	None
Indications	All Medically-accepted Indications.
Off-Label Uses	N/A
Part B Prerequisite	No

Formulary ID: 24449 version 13

Last Updated: 06/20/2024 Effective: 07/01/2024



ORSERDU

Products Affected

• ORSERDU

PA Criteria	Criteria Details
Exclusion Criteria	None
Required Medical Information	Diagnosis of advanced or metastatic, ER-positive, HER2-negative, ESR1-mutated, breast cancer in postmenopausal women or adult men after at least 1 line of endocrine therapy
Age Restrictions	18 years of age and older
Prescriber Restrictions	Prescribed by or in consultation with an oncologist
Coverage Duration	12 months
Other Criteria	None
Indications	All Medically-accepted Indications.
Off-Label Uses	N/A
Part B Prerequisite	No

Formulary ID: 24449 version 13

Last Updated: 06/20/2024 Effective: 07/01/2024



OSPHENA

Products Affected

OSPHENA

DA Cuitorio	Cuitouio Dotoila
PA Criteria	Criteria Details
Exclusion Criteria	Any of the following A.) Undiagnosed abnormal genital bleeding, B.) Known or suspected estrogen-dependent neoplasia, C.) Active deep vein thrombosis (DVT), pulmonary embolism (PE), or a history of these conditions, D.) Active arterial thromboembolic disease (e.g. stroke, myocardial infarction) or a history of these conditions, or E.) Pregnancy
Required Medical Information	Diagnosis of one of the following A.) Moderate to severe dyspareunia due to vulvar and vaginal atrophy associated with menopause, or B.) Moderate to severe vaginal dryness due to vulvar and vaginal atrophy associated with menopause
Age Restrictions	18 years of age and older
Prescriber Restrictions	None
Coverage Duration	12 months
Other Criteria	None
Indications	All Medically-accepted Indications.
Off-Label Uses	N/A
Part B Prerequisite	No

Formulary ID: 24449 version 13 Last Updated: 06/20/2024

Effective: 07/01/2024



OTEZLA

Products Affected

OTEZLA

PA Criteria	Criteria Details
Exclusion Criteria	None
Required Medical Information	Diagnosis of one of the following: A.) Active psoriatic arthritis and patient has trial and failure or intolerance or contraindication to two preferred products, (i.e. Humira, Enbrel, Rinvoq, Skyrizi, Stelara), B.) Moderate to severe plaque psoriasis, patient is a candidate for phototherapy or systemic therapy, and patient has trial and failure or intolerance or contraindication to two preferred products, (i.e. Humira, Enbrel, Skyrizi, Stelara), C.) Mild plaque psoriasis, patient is a candidate for phototherapy or systemic therapy, and patient has trial and failure or intolerance or contraindication to at least one topical psoriasis product (e.g., medium to high potency corticosteroid and/or vitamin D analog), or D.) Behcet's Disease and patient has active oral ulcers
Age Restrictions	6 years of age and older
Prescriber Restrictions	PsA: Prescribed by or in consultation with a dermatologist or rheumatologist. Plaque psoriasis: Prescribed by or in consultation with a dermatologist.
Coverage Duration	12 months
Other Criteria	None
Indications	All Medically-accepted Indications.
Off-Label Uses	N/A
Part B Prerequisite	No

Formulary ID: 24449 version 13 Last Updated: 06/20/2024

Effective: 07/01/2024



PANRETIN

Products Affected

• PANRETIN

PA Criteria	Criteria Details
Exclusion Criteria	None
Required Medical Information	Diagnosis of AIDS-related Kaposi's sarcoma and both of the following 1.) Used to treat cutaneous lesions, and 2.) Systemic anti-Kaposi's Sarcoma therapy is not indicated (e.g., patient does not have more than 10 new KS lesions in the prior month, symptomatic lymphedema, symptomatic pulmonary KS, or symptomatic visceral involvement)
Age Restrictions	18 years of age and older
Prescriber Restrictions	Prescribed by or in consultation with an oncologist or HIV specialist
Coverage Duration	12 months
Other Criteria	None
Indications	All Medically-accepted Indications.
Off-Label Uses	N/A
Part B Prerequisite	No

Formulary ID: 24449 version 13 Last Updated: 06/20/2024

Effective: 07/01/2024



PAZOPANIB

Products Affected

• pazopanib hcl

PA Criteria	Criteria Details
Exclusion Criteria	None
Required Medical Information	Diagnosis of one of the following A.) Advanced renal cell carcinoma, or B.) Advanced soft tissue sarcoma and patient received at least one prior chemotherapy
Age Restrictions	18 years of age and older
Prescriber Restrictions	Prescribed by or in consultation with an oncologist
Coverage Duration	12 months
Other Criteria	None
Indications	All Medically-accepted Indications.
Off-Label Uses	N/A
Part B Prerequisite	No

Formulary ID: 24449 version 13

Last Updated: 06/20/2024 Effective: 07/01/2024



PEGYLATED INTERFERON

Products Affected

- PEGASYS SUBCUTANEOUS SOLUTION 180 MCG/ML
- PEGASYS SUBCUTANEOUS SOLUTION PREFILLED SYRINGE

PA Criteria	Criteria Details
Exclusion Criteria	Any of the following A.) Autoimmune hepatitis, B.) Hepatic decompensation (Child-Pugh score greater than 6 (Class B and C) in cirrhotic patients before treatment, OR hepatic decompensation (Child-Pugh score greater than or equal to 6) in cirrhotic patients co-infected with hepatitis C and HIV before treatment, C.) Hypersensitivity reactions, including urticaria, bronchoconstriction, anaphylaxis, or Stevens-Johnson syndrome to alfa interferons or any component of the product, or D.) Pregnancy with concomitant ribavirin use
Required Medical Information	Diagnosis of one of the following A.) Chronic hepatitis B infection, or B.) Chronic hepatitis C and required criteria will be applied consistent with current AASLD-IDSA guidance with compensated liver disease
Age Restrictions	None
Prescriber Restrictions	Prescribed by or in consultation with a gastroenterologist, hepatologist, or infectious disease specialist
Coverage Duration	12 months
Other Criteria	None
Indications	All Medically-accepted Indications.
Off-Label Uses	N/A
Part B Prerequisite	No

Formulary ID: 24449 version 13 Last Updated: 06/20/2024

Effective: 07/01/2024



PEMAZYRE

Products Affected

PEMAZYRE

PA Criteria	Criteria Details
Exclusion Criteria	None
Required Medical Information	Diagnosis of one of the following A.) Previously treated, unresectable locally advanced or metastatic cholangiocarcinoma with confirmed fibroblast growth factor receptor 2 (FGFR2) fusion or other rearrangement as detected by an FDA-approved test or B.) Relapsed or refractory myeloid/lymphoid neoplasms with fibroblast growth factor receptor 1 rearrangement
Age Restrictions	18 years of age and older
Prescriber Restrictions	Prescribed by or in consultation with an oncologist, gastroenterologist, or hepatologist
Coverage Duration	12 months
Other Criteria	None
Indications	All Medically-accepted Indications.
Off-Label Uses	N/A
Part B Prerequisite	No

Formulary ID: 24449 version 13 Last Updated: 06/20/2024

Effective: 07/01/2024



PIQRAY

Products Affected

- PIQRAY (200 MG DAILY DOSE)
- PIQRAY (250 MG DAILY DOSE)

• PIQRAY (300 MG DAILY DOSE)

PA Criteria	Criteria Details
Exclusion Criteria	None
Required Medical Information	Diagnosis of hormone receptor (HR) positive, HER2-negative, PIK3CA-mutated, advanced or metastatic breast cancer and must meet all of the following 1.) Used in combination with fulvestrant, 2.) Disease has progressed on or after an endocrine-based regimen, and 3.) Patient is a male or postmenopausal female
Age Restrictions	18 years of age and older
Prescriber Restrictions	Prescribed by or in consultation with an oncologist
Coverage Duration	12 months
Other Criteria	None
Indications	All Medically-accepted Indications.
Off-Label Uses	N/A
Part B Prerequisite	No

Formulary ID: 24449 version 13

Last Updated: 06/20/2024 Effective: 07/01/2024



PIRFENIDONE

Products Affected

• pirfenidone

PA Criteria	Criteria Details
Exclusion Criteria	None
Required Medical Information	Diagnosis of idiopathic pulmonary fibrosis
Age Restrictions	18 years of age and older
Prescriber Restrictions	Prescribed by or in consultation with a pulmonologist
Coverage Duration	12 months
Other Criteria	None
Indications	All Medically-accepted Indications.
Off-Label Uses	N/A
Part B Prerequisite	No

Formulary ID: 24449 version 13 Last Updated: 06/20/2024

Effective: 07/01/2024



POMALYST

Products Affected

POMALYST

PA Criteria	Criteria Details
Exclusion Criteria	Pregnancy
Required Medical Information	Diagnosis of one of the following A.) AIDS-related Kaposi sarcoma and patient has failure on highly active antiretroviral therapy (HAART), B.) Kaposi sarcoma in HIV-negative adults, or C.) Multiple myeloma and in combination with dexamethasone in adults who have received at least 2 prior therapies (including lenalidomide and a proteasome inhibitor) and have demonstrated disease progression on or within 60 days of completion of the last therapy
Age Restrictions	18 years of age and older
Prescriber Restrictions	None
Coverage Duration	12 months
Other Criteria	None
Indications	All Medically-accepted Indications.
Off-Label Uses	N/A
Part B Prerequisite	No

Formulary ID: 24449 version 13 Last Updated: 06/20/2024

Effective: 07/01/2024



POSACONAZOLE

Products Affected

• posaconazole oral

PA Criteria	Criteria Details
Exclusion Criteria	Any of the following A.) Concomitant treatment with sirolimus, B.) Concomitant use of CYP3A4 substrates that prolong QT interval (pimozide, quinidine), C.) Concomitant use of HMG-CoA Reductase inhibitors primarily metabolized through CYP3A4, or D.) Concomitant use of ergot alkaloids
Required Medical Information	Diagnosis of one of the following A.) Oropharyngeal candidiasis, B.) Patient is severely immunocompromised and requires prophylaxis of invasive aspergillosis or candidiasis due to high risk of infection, or C.) Invasive aspergillosis
Age Restrictions	None
Prescriber Restrictions	None
Coverage Duration	12 weeks
Other Criteria	None
Indications	All Medically-accepted Indications.
Off-Label Uses	N/A
Part B Prerequisite	No

Formulary ID: 24449 version 13

Last Updated: 06/20/2024 Effective: 07/01/2024



POSACONAZOLE SUSPENSION

Products Affected

• NOXAFIL ORAL PACKET

• posaconazole oral

PA Criteria	Criteria Details
Exclusion Criteria	Any of the following A.) Concomitant treatment with sirolimus, B.) Concomitant use of CYP3A4 substrates that prolong QT interval (pimozide, quinidine), C.) Concomitant use of HMG-CoA Reductase inhibitors primarily metabolized through CYP3A4, or D.) Concomitant use of ergot alkaloids
Required Medical Information	Diagnosis of one of the following A.) Oropharyngeal candidiasis, B.) Patient is severely immunocompromised and requires prophylaxis of invasive aspergillosis or candidiasis due to high risk of infection, or C.) Invasive aspergillosis
Age Restrictions	None
Prescriber Restrictions	None
Coverage Duration	12 weeks
Other Criteria	None
Indications	All Medically-accepted Indications.
Off-Label Uses	N/A
Part B Prerequisite	No

Formulary ID: 24449 version 13 Last Updated: 06/20/2024

Effective: 07/01/2024



PREVYMIS

Products Affected

• PREVYMIS ORAL

PA Criteria	Criteria Details
Exclusion Criteria	Any of the following A.) Concomitant use with pimozide or ergot alkaloids (ergotamine, dihydroergotamine), B.) Concomitant use with pitavastatin or simvastatin when coadministered with cyclosporine
Required Medical Information	Diagnosis of one of the following A.) Prophylaxis of cytomegalovirus (CMV) infection and disease in adult CMV-seropositive recipients [R+] of an allogeneic hematopoietic stem cell transplant, or B.) Prophylaxis of CMV disease in adult kidney transplant recipients at high risk (Donor CMV seropositive/Recipient CMV seronegative [D+/R-])
Age Restrictions	18 years of age and older
Prescriber Restrictions	None
Coverage Duration	6 months
Other Criteria	None
Indications	All Medically-accepted Indications.
Off-Label Uses	N/A
Part B Prerequisite	No

Formulary ID: 24449 version 13 Last Updated: 06/20/2024

Effective: 07/01/2024



PROMACTA

Products Affected

• PROMACTA

PA Criteria	Criteria Details
Exclusion Criteria	None
Required Medical Information	Diagnosis of one of the following A.) Chronic idiopathic thrombocytopenic purpura (ITP), B.) Chronic hepatitis C infection associated thrombocytopenia, or C.) Severe aplastic anemia with insufficient response to immunosuppressive therapy or in combination with standard immunosuppressive therapy
Age Restrictions	None
Prescriber Restrictions	None
Coverage Duration	12 months
Other Criteria	None
Indications	All Medically-accepted Indications.
Off-Label Uses	N/A
Part B Prerequisite	No

Formulary ID: 24449 version 13

Last Updated: 06/20/2024 Effective: 07/01/2024



QINLOCK

Products Affected

• QINLOCK

PA Criteria	Criteria Details
Exclusion Criteria	None
Required Medical Information	Diagnosis of advanced gastrointestinal stromal tumor (GIST) and patient has received prior treatment with 3 or more kinase inhibitors, including imatinib
Age Restrictions	18 years of age and older
Prescriber Restrictions	None
Coverage Duration	12 months
Other Criteria	None
Indications	All Medically-accepted Indications.
Off-Label Uses	N/A
Part B Prerequisite	No

Formulary ID: 24449 version 13

Last Updated: 06/20/2024 Effective: 07/01/2024

 $H9306_24_DRS_09\text{-}001_OE_C$



QUININE SULFATE

Products Affected

• quinine sulfate oral

PA Criteria	Criteria Details
Exclusion Criteria	Any of the following A.) Prolongation of QT interval, B.) Glucose-6-phosphate dehydrogenase deficiency, C.) Myasthenia gravis, D.) Known hypersensitivity to mefloquine or quinidine, E.) Optic neuritis, F.) Diagnosis of Blackwater fever, G.) Use solely for treatment or prevention of nocturnal leg cramps
Required Medical Information	Diagnosis of one of the following A.) uncomplicated Plasmodium falciparum malaria, B.) uncomplicated Plasmodium vivax malaria, or C.) babesiosis
Age Restrictions	None
Prescriber Restrictions	None
Coverage Duration	1 month
Other Criteria	None
Indications	All Medically-accepted Indications.
Off-Label Uses	N/A
Part B Prerequisite	No

Formulary ID: 24449 version 13

Last Updated: 06/20/2024 Effective: 07/01/2024



RAVICTI

Products Affected

• RAVICTI

PA Criteria	Criteria Details
Exclusion Criteria	None
Required Medical Information	Diagnosis of urea cycle disorders
Age Restrictions	None
Prescriber Restrictions	None
Coverage Duration	12 months
Other Criteria	None
Indications	All Medically-accepted Indications.
Off-Label Uses	N/A
Part B Prerequisite	No

Formulary ID: 24449 version 13

Last Updated: 06/20/2024 Effective: 07/01/2024

 $H9306_24_DRS_09\text{-}001_OE_C$



REGRANEX

Products Affected

REGRANEX

PA Criteria	Criteria Details
Exclusion Criteria	Known neoplasm at the site of application
Required Medical Information	Diagnosis of lower extremity diabetic neuropathic ulcers that extend into the subcutaneous tissue or beyond and have an adequate blood supply
Age Restrictions	16 years of age and older
Prescriber Restrictions	None
Coverage Duration	12 months
Other Criteria	None
Indications	All Medically-accepted Indications.
Off-Label Uses	N/A
Part B Prerequisite	No

Formulary ID: 24449 version 13

Last Updated: 06/20/2024 Effective: 07/01/2024



REPATHA

Products Affected

REPATHA

- REPATHA SURECLICK
- REPATHA PUSHTRONEX SYSTEM

PA Criteria	Criteria Details
Exclusion Criteria	None
Required Medical Information	Diagnosis of one of the following A.) primary hyperlipidemia including heterozygous familial hypercholesterolemia (HeFH), B.) homozygous familial hypercholesterolemia, C.) established cardiovascular disease and myocardial infarction prophylaxis, stroke prophylaxis, or coronary revascularization prophylaxis is required, or D.) clinical atherosclerotic cardiovascular disease (CVD) as defined as one of the following 1.) acute coronary syndrome, 2.) history of myocardial infarction, 3.) stable/unstable angina, 4.) coronary or other arterial revascularization, 5.) stroke, 6.) transient ischemic stroke (TIA), or 7.) peripheral arterial disease presumed to be atherosclerotic region
Age Restrictions	10 years of age and older
Prescriber Restrictions	None
Coverage Duration	12 months
Other Criteria	None
Indications	All Medically-accepted Indications.
Off-Label Uses	N/A
Part B Prerequisite	No

Formulary ID: 24449 version 13 Last Updated: 06/20/2024

Effective: 07/01/2024



RETEVMO

Products Affected

RETEVMO

PA Criteria	Criteria Details
Exclusion Criteria	None
Required Medical Information	Diagnosis of one of the following A.) Advanced or metastatic RET-mutant medullary thyroid cancer (MTC) in patients who require systemic therapy, B.) Metastatic RET fusion-positive non-small cell lung cancer (NSCLC), C.) Advanced or metastatic RET fusion-positive thyroid cancer in patients who require systemic therapy and are refractory to radioactive iodine, if appropriate, or D.) Locally advanced or metastatic solid tumors with a RET gene fusion that have progressed on or following prior systemic treatment or who have no satisfactory alternative treatment options
Age Restrictions	None
Prescriber Restrictions	Prescribed by or in consultation with an oncologist
Coverage Duration	12 months
Other Criteria	None
Indications	All Medically-accepted Indications.
Off-Label Uses	N/A
Part B Prerequisite	No

Formulary ID: 24449 version 13 Last Updated: 06/20/2024

Effective: 07/01/2024



REZLIDHIA

Products Affected

• REZLIDHIA

PA Criteria	Criteria Details
Exclusion Criteria	None
Required Medical Information	Diagnosis of relapsed or refractory acute myeloid leukemia (AML) with a susceptible IDH1 mutation as detected by an FDA-approved test
Age Restrictions	18 years of age and older
Prescriber Restrictions	Prescribed by or in consultation with an oncologist
Coverage Duration	12 months
Other Criteria	None
Indications	All Medically-accepted Indications.
Off-Label Uses	N/A
Part B Prerequisite	No

Formulary ID: 24449 version 13

Last Updated: 06/20/2024 Effective: 07/01/2024



REZUROCK

Products Affected

REZUROCK

PA Criteria	Criteria Details
Exclusion Criteria	None
Required Medical Information	Diagnosis of chronic graft-vs-host disease and patient has failed at least 2 prior lines of systemic therapy.
Age Restrictions	12 years of age and older
Prescriber Restrictions	None
Coverage Duration	12 months
Other Criteria	None
Indications	All Medically-accepted Indications.
Off-Label Uses	N/A
Part B Prerequisite	No

Formulary ID: 24449 version 13 Last Updated: 06/20/2024

Effective: 07/01/2024



RILUZOLE

Products Affected

• riluzole

PA Criteria	Criteria Details
Exclusion Criteria	None
Required Medical Information	Diagnosis of amyotrophic lateral sclerosis (ALS)
Age Restrictions	18 years of age and older
Prescriber Restrictions	None
Coverage Duration	12 months
Other Criteria	None
Indications	All Medically-accepted Indications.
Off-Label Uses	N/A
Part B Prerequisite	No

Formulary ID: 24449 version 13 Last Updated: 06/20/2024

Effective: 07/01/2024



RINVOQ

Products Affected

• RINVOQ

PA Criteria	Criteria Details
Exclusion Criteria	None
Required Medical Information	Diagnosis of one of the following A.) Moderate to severe rheumatoid arthritis, B.) Active psoriatic arthritis, C.) Moderate to severe atopic dermatitis and patient has trial/failure, contraindication, or intolerance to two of the following 1.) Topical corticosteroid and/or 2.) Topical calcineurin inhibitor, D.) Moderately to severely active ulcerative colitis who have had an inadequate response or intolerance to one or more tumor necrosis factor blockers, E.) Active ankylosing spondylitis who have had an inadequate response or intolerance to one or more tumor necrosis factor blockers, F.) Active nonradiographic axial spondyloarthritis with objective signs of inflammation who have had an inadequate response or intolerance to tumor necrosis factor blocker therapy, or G.) Moderate to severe active Crohn's disease who have had an inadequate response or intolerance to tumor necrosis factor blocker therapy
Age Restrictions	None
Prescriber Restrictions	None
Coverage Duration	12 months
Other Criteria	Screening for latent tuberculosis infection is required prior to initiation of treatment
Indications	All Medically-accepted Indications.
Off-Label Uses	N/A
Part B Prerequisite	No

Formulary ID: 24449 version 13 Last Updated: 06/20/2024

Effective: 07/01/2024



RIVFLOZA

Products Affected

• RIVFLOZA

PA Criteria	Criteria Details
Exclusion Criteria	None
Required Medical Information	Diagnosis of Primary hyperoxaluria type 1 and the patient has relatively preserved kidney function (eGFR is greater than or equal to 30mL/min/1.73m(2)
Age Restrictions	9 years of age and older
Prescriber Restrictions	None
Coverage Duration	12 months
Other Criteria	None
Indications	All Medically-accepted Indications.
Off-Label Uses	N/A
Part B Prerequisite	No

Formulary ID: 24449 version 13 Last Updated: 06/20/2024

Effective: 07/01/2024



ROZLYTREK

Products Affected

ROZLYTREK

PA Criteria	Criteria Details
Exclusion Criteria	None
Required Medical Information	Diagnosis of one of the following A.) ROS1-positive metastatic non-small cell lung cancer (NSCLC), or B.) Solid tumors that have a neurotrophic tyrosine receptor kinase (NTRK) gene fusion without a known acquired resistance mutation, are metastatic or where surgical resection is likely to result in severe morbidity, and have either progressed following treatment or have no satisfactory alternative therapy
Age Restrictions	None
Prescriber Restrictions	Prescribed by or in consultation with an oncologist
Coverage Duration	12 months
Other Criteria	None
Indications	All Medically-accepted Indications.
Off-Label Uses	N/A
Part B Prerequisite	No

Formulary ID: 24449 version 13 Last Updated: 06/20/2024

Effective: 07/01/2024



RUBRACA

Products Affected

• RUBRACA

PA Criteria	Criteria Details
Exclusion Criteria	None
Required Medical Information	Diagnosis of one of the following A.) Recurrent epithelial ovarian, fallopian tube, or primary peritoneal cancer, used as maintenance treatment, and patient is in complete or partial response to platinum-based chemotherapy, or B.) Deleterious BRCA mutation (germline and/or somatic)-associated metastatic castration-resistant prostate cancer and patient has been treated with androgen receptor-directed therapy and a taxane-based chemotherapy
Age Restrictions	18 years of age and older
Prescriber Restrictions	Prescribed by or in consultation with an oncologist or hematologist
Coverage Duration	12 months
Other Criteria	None
Indications	All Medically-accepted Indications.
Off-Label Uses	N/A
Part B Prerequisite	No

Formulary ID: 24449 version 13

Last Updated: 06/20/2024 Effective: 07/01/2024



RYDAPT

Products Affected

• RYDAPT

PA Criteria	Criteria Details
Exclusion Criteria	None
Required Medical Information	Diagnosis of one of the following A.) treatment naive FLT3 mutation-positive acute myelogenous leukemia (AML) and must be used in combination with standard cytarabine and daunorubicin induction and cytarabine consolidation therapy, or B.) systemic mastocytosis or mast cell leukemia
Age Restrictions	18 years of age and older
Prescriber Restrictions	Prescribed by or in consultation with an oncologist or hematologist
Coverage Duration	12 months
Other Criteria	None
Indications	All Medically-accepted Indications.
Off-Label Uses	N/A
Part B Prerequisite	No

Formulary ID: 24449 version 13 Last Updated: 06/20/2024

Effective: 07/01/2024



SAPROPTERIN

Products Affected

• sapropterin dihydrochloride oral packet • sapropterin dihydrochloride oral tablet

PA Criteria	Criteria Details
Exclusion Criteria	None
Required Medical Information	Diagnosis of hyperphenylalaninemia (HPA) caused by tetrahydrobiopterin (BH4)-responsive phenylketonuria (PKU)
Age Restrictions	None
Prescriber Restrictions	None
Coverage Duration	Initial: 2 months, Renewal: 12 months
Other Criteria	None
Indications	All Medically-accepted Indications.
Off-Label Uses	N/A
Part B Prerequisite	No

Formulary ID: 24449 version 13 Last Updated: 06/20/2024

Effective: 07/01/2024



SCEMBLIX

Products Affected

• SCEMBLIX

PA Criteria	Criteria Details
Exclusion Criteria	None
Required Medical Information	Diagnosis of one of the following A.) Philadelphia chromosome-positive chronic myeloid leukemia (Ph+ CML) in chronic phase (CP), previously treated with two or more tyrosine kinase inhibitors (TKIs), or B.) Philadelphia chromosome-positive chronic myeloid leukemia (Ph+ CML) in chronic phase (CP) with the T315I mutation
Age Restrictions	18 years of age and older
Prescriber Restrictions	Prescribed by or in consultation with an oncologist or hematologist
Coverage Duration	12 months
Other Criteria	None
Indications	All Medically-accepted Indications.
Off-Label Uses	N/A
Part B Prerequisite	No

Formulary ID: 24449 version 13

Last Updated: 06/20/2024 Effective: 07/01/2024



SIGNIFOR

Products Affected

• SIGNIFOR

PA Criteria	Criteria Details
Exclusion Criteria	None
Required Medical Information	Diagnosis of Cushing disease and patient has had inadequate response to or is not a candidate for surgery. For renewal: Documentation of a clinically meaningful reduction in 24-hour urinary free cortisol (UFC) levels or improvement in signs or symptoms of the disease
Age Restrictions	18 years of age and older
Prescriber Restrictions	None
Coverage Duration	Initial: 6 months, Renewal: 12 months
Other Criteria	None
Indications	All Medically-accepted Indications.
Off-Label Uses	N/A
Part B Prerequisite	No

Formulary ID: 24449 version 13 Last Updated: 06/20/2024

Effective: 07/01/2024



SILDENAFIL

Products Affected

• sildenafil citrate oral tablet 20 mg

PA Criteria	Criteria Details
Exclusion Criteria	Any of the following A.) Nitrate therapy, including intermittent use, B.) Concomitant use with riocguat or other guanylate cyclase stimulators, C.) Concomitant use with HIV protease inhibitors or elvitegravir/cobicistat/tenofovir/emtricitabine
Required Medical Information	Diagnosis of pulmonary arterial hypertension (WHO Group I), confirmed by right heart catheterization or Doppler echocardiogram if patient is unable to undergo a right heart catheterization (e.g., patient is frail, elderly, etc.)
Age Restrictions	None
Prescriber Restrictions	Prescribed by or in consultation with a cardiologist or pulmonologist
Coverage Duration	12 months
Other Criteria	None
Indications	All Medically-accepted Indications.
Off-Label Uses	N/A
Part B Prerequisite	No

Formulary ID: 24449 version 13 Last Updated: 06/20/2024

Effective: 07/01/2024



SIRTURO

Products Affected

• SIRTURO

PA Criteria	Criteria Details
Exclusion Criteria	None
Required Medical Information	Must meet all of the following 1.) Diagnosis of pulmonary multidrug resistant tuberculosis (MDR-TB) and 2.) Used in combination with at least 3 other antibiotics for the treatment of pulmonary multi-drug resistant tuberculosis
Age Restrictions	5 years of age and older
Prescriber Restrictions	Prescribed by or in consultation with an infectious disease specialist
Coverage Duration	24 weeks
Other Criteria	None
Indications	All Medically-accepted Indications.
Off-Label Uses	N/A
Part B Prerequisite	No

Formulary ID: 24449 version 13 Last Updated: 06/20/2024

Effective: 07/01/2024



SKYRIZI

Products Affected

• SKYRIZI PEN

• SKYRIZI SUBCUTANEOUS

PA Criteria	Criteria Details
Exclusion Criteria	None
Required Medical Information	Diagnosis of one of the following A.) Moderate to severe plaque psoriasis and patient is a candidate for systemic therapy or phototherapy, B.) Active psoriatic arthritis, or C.) Moderately to severely active Crohn's disease
Age Restrictions	None
Prescriber Restrictions	None
Coverage Duration	12 months
Other Criteria	Screening for latent tuberculosis infection is required prior to initiation of treatment
Indications	All Medically-accepted Indications.
Off-Label Uses	N/A
Part B Prerequisite	No

Formulary ID: 24449 version 13 Last Updated: 06/20/2024

Effective: 07/01/2024



SODIUM OXYBATE

Products Affected

• sodium oxybate

PA Criteria	Criteria Details
Exclusion Criteria	Any of the following A.) Concomitant treatment with sedative hypnotic agents, B.) Succinic semialdehyde dehydrogenase deficiency
Required Medical Information	Diagnosis of one of the following A.) Narcolepsy with excessive daytime drowsiness and has trial of/or contraindication to a central nervous system (CNS) stimulant drug (e.g., amphetamine, dextroamphetamine, methylphenidate) or a CNS wakefulness promoting drug (e.g., armodafinil, modafinil), or B.) Cataplexy and narcolepsy
Age Restrictions	7 years of age and older
Prescriber Restrictions	None
Coverage Duration	12 months
Other Criteria	None
Indications	All Medically-accepted Indications.
Off-Label Uses	N/A
Part B Prerequisite	No

Formulary ID: 24449 version 13 Last Updated: 06/20/2024

Effective: 07/01/2024



SOHONOS

Products Affected

SOHONOS

PA Criteria	Criteria Details
Exclusion Criteria	Pregnancy
Required Medical Information	Diagnosis of fibrodysplasia ossificans progressiva (FOP)
Age Restrictions	8 years and older for females and 10 years and older for males
Prescriber Restrictions	None
Coverage Duration	6 months
Other Criteria	None
Indications	All Medically-accepted Indications.
Off-Label Uses	N/A
Part B Prerequisite	No

Formulary ID: 24449 version 13 Last Updated: 06/20/2024

Effective: 07/01/2024



SOLTAMOX

Products Affected

SOLTAMOX

PA Criteria	Criteria Details
Exclusion Criteria	Any of the following A.) Concomitant coumarin-type anticoagulant therapy, B.) history of thromboembolic disease such as DVT or PE
Required Medical Information	Diagnosis of breast cancer and documentation of inability to swallow tablet formulation
Age Restrictions	18 years of age and older
Prescriber Restrictions	Prescribed by or in consultation with an oncologist
Coverage Duration	12 months
Other Criteria	None
Indications	All Medically-accepted Indications.
Off-Label Uses	N/A
Part B Prerequisite	No

Formulary ID: 24449 version 13

Last Updated: 06/20/2024 Effective: 07/01/2024

 $H9306_24_DRS_09\text{-}001_OE_C$



SOMAVERT

Products Affected

SOMAVERT

PA Criteria	Criteria Details
Exclusion Criteria	None
Required Medical Information	Diagnosis of acromegaly confirmed by high pretreatment insulin-like growth factor-1 (IGF-1) level for age and/or gender based on the laboratory reference range and patient has had an inadequate response to or is ineligible for surgery or radiation therapy
Age Restrictions	18 years of age and older
Prescriber Restrictions	Prescribed by or in consultation with an endocrinologist
Coverage Duration	12 months
Other Criteria	None
Indications	All Medically-accepted Indications.
Off-Label Uses	N/A
Part B Prerequisite	No

Formulary ID: 24449 version 13

Last Updated: 06/20/2024 Effective: 07/01/2024



SORAFENIB

Products Affected

• sorafenib tosylate

PA Criteria	Criteria Details
Exclusion Criteria	Squamous cell lung cancer being treated with carboplatin and paclitaxel
Required Medical Information	Diagnosis of one of the following A.) Advanced renal cell carcinoma, B.) Locally recurrent or metastatic, progressive, differentiated thyroid carcinoma that is refractory to radioactive iodine treatment, or C.) Unresectable hepatocellular carcinoma
Age Restrictions	18 years of age and older
Prescriber Restrictions	None
Coverage Duration	12 months
Other Criteria	None
Indications	All Medically-accepted Indications.
Off-Label Uses	N/A
Part B Prerequisite	No

Formulary ID: 24449 version 13

Last Updated: 06/20/2024 Effective: 07/01/2024

 $H9306_24_DRS_09\text{-}001_OE_C$



SPRYCEL

Products Affected

• SPRYCEL

PA Criteria	Criteria Details
Exclusion Criteria	None
Required Medical Information	Diagnosis of one of the following A.) Newly diagnosed Philadelphia chromosome-positive (Ph+) chronic myeloid leukemia (CML) in chronic phase, B.) Chronic, accelerated, or myeloid or lymphoid blast phase Ph+ CML with resistance or intolerance to prior therapy, C.) Ph+ acute lymphoblastic leukemia (ALL) with resistance or intolerance to prior therapy, or D.) Newly diagnosed Ph+ ALL in combination with chemotherapy
Age Restrictions	1 years of age and older
Prescriber Restrictions	Prescribed by or in consultation with an oncologist or hematologist
Coverage Duration	12 months
Other Criteria	None
Indications	All Medically-accepted Indications.
Off-Label Uses	N/A
Part B Prerequisite	No

Formulary ID: 24449 version 13

Last Updated: 06/20/2024 Effective: 07/01/2024



STELARA

Products Affected

- STELARA SUBCUTANEOUS SOLUTION 45 MG/0.5ML
- STELARA SUBCUTANEOUS SOLUTION PREFILLED SYRINGE

PA Criteria	Criteria Details
Exclusion Criteria	None
Required Medical Information	Diagnosis of one of the following A.) Moderate to severely active Crohn disease, B.) Moderate to severe plaque psoriasis, C.) Active psoriatic arthritis, or D.) Moderate to severe active ulcerative colitis
Age Restrictions	None
Prescriber Restrictions	None
Coverage Duration	12 months
Other Criteria	Screening for latent tuberculosis infection is required prior to initiation of treatment
Indications	All Medically-accepted Indications.
Off-Label Uses	N/A
Part B Prerequisite	No

Formulary ID: 24449 version 13 Last Updated: 06/20/2024

Effective: 07/01/2024



STIVARGA

Products Affected

• STIVARGA

PA Criteria	Criteria Details
ra Criteria	Criteria Details
Exclusion Criteria	None
Required Medical Information	Diagnosis of one of the following A.) Metastatic colorectal cancer in patients previously treated with fluoropyrimidine, oxaliplatin, and irinotecan containing chemotherapy, anti-VEGF therapy, and if RAS wild type, anti-EGFR therapy, B.) Liver carcinoma in patients previously treated with sorafenib, or C.) Locally advanced, unresectable or metastatic gastrointestinal stromal tumor (GIST) after treatment with imatinib and sunitinib
Age Restrictions	18 years of age and older
Prescriber Restrictions	None
Coverage Duration	12 months
Other Criteria	None
Indications	All Medically-accepted Indications.
Off-Label Uses	N/A
Part B Prerequisite	No

Formulary ID: 24449 version 13

Last Updated: 06/20/2024 Effective: 07/01/2024



SUNITINIB

Products Affected

• sunitinib malate

PA Criteria	Criteria Details
Exclusion Criteria	None
Required Medical Information	Diagnosis of one of the following A.) Gastrointestinal stromal tumor after disease progression on or intolerance to imatinib, B.) Pancreatic neuroendocrine tumors in a patient with unresectable locally advanced or metastatic disease, C.) Advanced renal cell carcinoma, or D.) Renal cell carcinoma and used as adjuvant therapy following nephrectomy in patients who are at high risk for recurrence
Age Restrictions	18 years of age and older
Prescriber Restrictions	None
Coverage Duration	12 months
Other Criteria	None
Indications	All Medically-accepted Indications.
Off-Label Uses	N/A
Part B Prerequisite	No

Formulary ID: 24449 version 13

Last Updated: 06/20/2024 Effective: 07/01/2024



SYMDEKO

Products Affected

• SYMDEKO

PA Criteria	Criteria Details
Exclusion Criteria	None
Required Medical Information	Diagnosis of cystic fibrosis (CF) and must meet one of the following 1.) Patient is homozygous for the F508del mutation, or 2.) Patient has at least one mutation in the cystic fibrosis transmembrane conductance regulator (CFTR) gene that is responsive to tezacaftor/ivacaftor verified by an FDA-cleared CF mutation test
Age Restrictions	6 years of age and older
Prescriber Restrictions	Prescribed by or in consultation with a pulmonologist or prescribing practitioner is from a CF center accredited by the Cystic Fibrosis Foundation
Coverage Duration	Initial: 6 months, Renewal: 12 months
Other Criteria	None
Indications	All Medically-accepted Indications.
Off-Label Uses	N/A
Part B Prerequisite	No

Formulary ID: 24449 version 13 Last Updated: 06/20/2024

Effective: 07/01/2024



SYMLIN

Products Affected

- SYMLINPEN 120 SUBCUTANEOUS SOLUTION PEN-INJECTOR
- SYMLINPEN 60 SUBCUTANEOUS SOLUTION PEN-INJECTOR

PA Criteria	Criteria Details
Exclusion Criteria	Any of the following A.) Confirmed diagnosis of gastroparesis, B.) Hypoglycemia unawareness
Required Medical Information	Diagnosis of type 1 or type 2 diabetes mellitus and patient uses mealtime insulin therapy and has failed to achieve desired glucose control
Age Restrictions	18 years of age and older
Prescriber Restrictions	None
Coverage Duration	12 months
Other Criteria	None
Indications	All Medically-accepted Indications.
Off-Label Uses	N/A
Part B Prerequisite	No

Formulary ID: 24449 version 13

Last Updated: 06/20/2024 Effective: 07/01/2024



SYNAREL

Products Affected

• SYNAREL

PA Criteria	Criteria Details
Exclusion Criteria	Any of the following A.) pregnancy, B.) breastfeeding, C.) undiagnosed abnormal vaginal bleeding
Required Medical Information	Diagnosis of one of the following A.) Central precocious puberty, or B.) Endometriosis
Age Restrictions	None
Prescriber Restrictions	None
Coverage Duration	12 months
Other Criteria	None
Indications	All Medically-accepted Indications.
Off-Label Uses	N/A
Part B Prerequisite	No

Formulary ID: 24449 version 13

Last Updated: 06/20/2024 Effective: 07/01/2024



TABLOID

Products Affected

• TABLOID

PA Criteria	Criteria Details
Exclusion Criteria	None
Required Medical Information	Diagnosis of acute myeloid leukemia (induction and consolidation therapy only)
Age Restrictions	None
Prescriber Restrictions	Prescribed by or in consultation with an oncologist or hematologist
Coverage Duration	12 months
Other Criteria	None
Indications	All Medically-accepted Indications.
Off-Label Uses	N/A
Part B Prerequisite	No

Formulary ID: 24449 version 13

Last Updated: 06/20/2024 Effective: 07/01/2024



TABRECTA

Products Affected

• TABRECTA

PA Criteria	Criteria Details
Exclusion Criteria	None
Required Medical Information	Diagnosis of metastatic non-small cell lung cancer (NSCLC) in patients whose tumors have a mutation that leads to mesenchymal-epithelial transition (MET) exon 14 skipping as detected by an FDA-approved test
Age Restrictions	18 years of age and older
Prescriber Restrictions	Prescribed by or in consultation with an oncologist
Coverage Duration	12 months
Other Criteria	None
Indications	All Medically-accepted Indications.
Off-Label Uses	N/A
Part B Prerequisite	No

Formulary ID: 24449 version 13 Last Updated: 06/20/2024

Effective: 07/01/2024



TAFINLAR

Products Affected

TAFINLAR

PA Criteria	Criteria Details
Exclusion Criteria	None
Required Medical Information	Diagnosis of one of the following A.) Locally advanced or metastatic anaplastic thyroid carcinoma with BRAF V600E mutation, in combination with trametinib and no satisfactory locoregional treatment options, B.) Metastatic non-small cell lung cancer with BRAF V600E mutation, in combination with trametinib OR in patients previously treated as monotherapy, C.) Unresectable or metastatic malignant melanoma with BRAF V600E or V600K mutation, D.) Unresectable or metastatic solid tumors with BRAF V600E mutation, in combination with trametinib, and have progressed following prior treatment and have no satisfactory alternative treatment options, or E.) Low-grade glioma with a BRAF V600E mutation and require systemic therapy, in combination with trametinib
Age Restrictions	None
Prescriber Restrictions	Prescribed by or in consultation with an oncologist
Coverage Duration	12 months
Other Criteria	None
Indications	All Medically-accepted Indications.
Off-Label Uses	N/A
Part B Prerequisite	No

Formulary ID: 24449 version 13 Last Updated: 06/20/2024

Effective: 07/01/2024



TAGRISSO

Products Affected

TAGRISSO

PA Criteria	Criteria Details
Exclusion Criteria	None
Required Medical Information	Diagnosis of one of the following A.) Metastatic non-small cell lung cancer (NSCLC) with EGFR exon 19 deletion or exon 21 L858R mutation and used as first line therapy, B.) Metastatic non-small cell lung cancer with T790M EGFR mutation (as confirmed by an FDA-approved test) AND whose disease has progressed on or after EGFR tyrosine kinase inhibitor therapy, C.) Non-small cell lung cancer (NSCLC) with tumor epidermal growth factor receptor (EGFR) exon 19 deletions or exon 21 L858R mutations (as confirmed by an FDA-approved test) AND patient requires adjuvant therapy after tumor resection, or D.) First-line treatment of adult patients with locally advanced or metastatic non-small cell lung cancer whose tumors have epidermal growth factor receptor (EGFR) exon 19 deletions or exon 21 L858R mutations, as detected by an FDA-approved test, in combination with pemetrexed and platinum-based chemotherapy
Age Restrictions	18 years of age and older
Prescriber Restrictions	Prescribed by or in consultation with an oncologist
Coverage Duration	12 months
Other Criteria	None
Indications	All Medically-accepted Indications.
Off-Label Uses	N/A
Part B Prerequisite	No

Formulary ID: 24449 version 13 Last Updated: 06/20/2024

Effective: 07/01/2024



TAKHZYRO

Products Affected

TAKHZYRO

PA Criteria	Criteria Details
Exclusion Criteria	None
Required Medical Information	Diagnosis of one of the following and used as routine prophylaxis A.) Hereditary angioedema (HAE) with C1 inhibitor deficiency (Type 1) confirmed by laboratory testing, or B.) HAE with C1 inhibitor dysfunction (Type 2) confirmed by laboratory testing, or C.) HAE with normal C1 inhibitor (Type 3) confirmed by laboratory testing and one of the following 1.) Positive test for an F12, angiopoietin-1, or plasminogen gene mutation, or 2.) Family history of angioedema and the angioedema was refractory to a trial of an antihistamine for at least one month
Age Restrictions	None
Prescriber Restrictions	Prescribed by or in consultation with a hematologist, immunologist, or allergist
Coverage Duration	12 months
Other Criteria	None
Indications	All Medically-accepted Indications.
Off-Label Uses	N/A
Part B Prerequisite	No

Formulary ID: 24449 version 13 Last Updated: 06/20/2024

Effective: 07/01/2024



TALZENNA

Products Affected

TALZENNA

PA Criteria	Criteria Details
Exclusion Criteria	None
Required Medical Information	Diagnosis of one of the following A.) Deleterious or suspected deleterious germline breast cancer susceptibility gene (BRCA)-mutated (gBRCAm), human epidermal growth factor receptor 2 (HER2)-negative locally advanced or metastatic breast cancer, or B.) Homologous recombination repair gene-mutated metastatic castration-resistant prostate cancer in combination with enzulatamide
Age Restrictions	18 years of age and older
Prescriber Restrictions	Prescribed by or in consultation with an oncologist
Coverage Duration	12 months
Other Criteria	None
Indications	All Medically-accepted Indications.
Off-Label Uses	N/A
Part B Prerequisite	No

Formulary ID: 24449 version 13 Last Updated: 06/20/2024

Effective: 07/01/2024



TASIGNA

Products Affected

• TASIGNA

PA Criteria	Criteria Details
Exclusion Criteria	Any of the following A.) Long QT syndrome, B.) Uncorrected hypokalemia, C.) Uncorrected hypomagnesemia
Required Medical Information	Diagnosis of one of the following A.) Newly diagnosed chronic phase Philadelphia chromosome-positive chronic myelogenous leukemia (CML), B.) Chronic phase or accelerated phase Philadelphia chromosome-positive CML in a patient with resistance or intolerance to prior therapy that included imatinib, or C.) Chronic phase Philadelphia chromosome-positive CML in a patient with resistance or intolerance to prior tyrosine-kinase inhibitor therapy
Age Restrictions	1 years of age and older
Prescriber Restrictions	Prescribed by or in consultation with an oncologist or hematologist
Coverage Duration	12 months
Other Criteria	None
Indications	All Medically-accepted Indications.
Off-Label Uses	N/A
Part B Prerequisite	No

Formulary ID: 24449 version 13 Last Updated: 06/20/2024

Effective: 07/01/2024



TAVNEOS

Products Affected

TAVNEOS

PA Criteria	Criteria Details
Exclusion Criteria	None
Required Medical Information	Diagnosis of severe active anti-neutrophil cytoplasmic autoantibody (ANCA)-associated vasculitis (granulomatosis with polyangiitis [GPA] and microscopic polyangiitis [MPA]) and both of the following apply 1.) Used as adjunctive treatment, and 2.) Used in combination with standard therapy including glucocorticoids
Age Restrictions	18 years of age and older
Prescriber Restrictions	None
Coverage Duration	12 months
Other Criteria	None
Indications	All Medically-accepted Indications.
Off-Label Uses	N/A
Part B Prerequisite	No

Formulary ID: 24449 version 13 Last Updated: 06/20/2024

Effective: 07/01/2024



TAZAROTENE

Products Affected

tazarotene external creamtazarotene external gel

• TAZORAC EXTERNAL CREAM 0.05 %

• iuzuroiene exiernai gei	
PA Criteria	Criteria Details
Exclusion Criteria	Pregnancy
Required Medical Information	Diagnosis of one of the following A.) Acne vulgaris and patient has trial with at least one generic topical acne product, or B.) Stable moderate to severe plaque psoriasis with 20% or less body surface area involvement and patient has trial with at least one other topical psoriasis product (e.g., medium to high potency corticosteroid and/or vitamin D analogs)
Age Restrictions	12 years of age and older
Prescriber Restrictions	None
Coverage Duration	12 months
Other Criteria	None
Indications	All Medically-accepted Indications.
Off-Label Uses	N/A
Part B Prerequisite	No

Formulary ID: 24449 version 13 Last Updated: 06/20/2024

Effective: 07/01/2024



TAZVERIK

Products Affected

TAZVERIK

PA Criteria	Criteria Details
Exclusion Criteria	None
Required Medical Information	Diagnosis of one of the following A.) Metastatic or locally advanced epithelioid sarcoma in patients not eligible for complete resection, B.) Relapsed or refractory follicular lymphoma in patients whose tumors are positive for an EZH2 mutation as detected by an FDA-approved test and who have received at least 2 prior systemic therapies, or C.) Relapsed or refractory follicular lymphoma in patients who have no satisfactory alternative treatment options
Age Restrictions	16 years of age and older
Prescriber Restrictions	Prescribed by or in consultation with an oncologist
Coverage Duration	12 months
Other Criteria	None
Indications	All Medically-accepted Indications.
Off-Label Uses	N/A
Part B Prerequisite	No

Formulary ID: 24449 version 13

Last Updated: 06/20/2024 Effective: 07/01/2024



TEGSEDI

Products Affected

• TEGSEDI

PA Criteria	Criteria Details
Exclusion Criteria	Any of the following A.) Platelet count less than 100,000 per microliter, B.) Urinary protein to creatinine ratio (UPCR) of 1000 mg/g or higher
Required Medical Information	Diagnosis of Polyneuropathy of hereditary transthyretin-mediated amyloidosis
Age Restrictions	18 years of age and older
Prescriber Restrictions	None
Coverage Duration	12 months
Other Criteria	None
Indications	All Medically-accepted Indications.
Off-Label Uses	N/A
Part B Prerequisite	No

Formulary ID: 24449 version 13 Last Updated: 06/20/2024

Effective: 07/01/2024



TEPMETKO

Products Affected

TEPMETKO

PA Criteria	Criteria Details
Exclusion Criteria	None
Required Medical Information	Diagnosis of metastatic non-small cell lung cancer (NSCLC) with mesenchymal-epithelial transition (MET) exon 14 skipping alterations
Age Restrictions	18 years of age and older
Prescriber Restrictions	Prescribed by or in consultation with an oncologist
Coverage Duration	12 months
Other Criteria	None
Indications	All Medically-accepted Indications.
Off-Label Uses	N/A
Part B Prerequisite	No

Formulary ID: 24449 version 13

Last Updated: 06/20/2024 Effective: 07/01/2024



TERIPARATIDE

Products Affected

• teriparatide (recombinant) subcutaneous solution pen-injector 620 mcg/2.48ml

PA Criteria	Criteria Details
Exclusion Criteria	None
Required Medical Information	Diagnosis of one of the following A.) Osteoporosis in postmenopausal female patient with high risk for fracture and patient has contraindication or has tried/had inadequate response to a bisphosphonate or Tymlos, B.) Primary or hypogonadal osteoporosis in male patient with high risk for fracture and patient has contraindication or has tried/had inadequate response to a bisphosphonate, or C.) Osteoporosis due to associated sustained systemic glucocorticoid therapy in patient with high risk for fracture and patient has contraindication or has tried/had inadequate response to a bisphosphonate
Age Restrictions	18 years of age and older
Prescriber Restrictions	None
Coverage Duration	Initial: 12 months, Renewal: 12 months (Maximum 24 month treatment per patient lifetime)
Other Criteria	None
Indications	All Medically-accepted Indications.
Off-Label Uses	N/A
Part B Prerequisite	No

Formulary ID: 24449 version 13 Last Updated: 06/20/2024

Effective: 07/01/2024

 $H9306_24_DRS_09\text{-}001_OE_C$



TETRABENAZINE

Products Affected

• tetrabenazine

PA Criteria	Criteria Details
Exclusion Criteria	Any of the following A.) Actively suicidal, B.) Untreated or inadequately treated depression, C.) Impaired hepatic function, D.) Concomitant use of monoamine oxidase inhibitors, E.) Concomitant use of reserpine or within 20 days of discontinuing reserpine
Required Medical Information	Diagnosis of chorea associated with Huntington's disease
Age Restrictions	None
Prescriber Restrictions	None
Coverage Duration	12 months
Other Criteria	None
Indications	All Medically-accepted Indications.
Off-Label Uses	N/A
Part B Prerequisite	No

Formulary ID: 24449 version 13

Last Updated: 06/20/2024 Effective: 07/01/2024



THALOMID

Products Affected

• THALOMID

PA Criteria	Criteria Details
Exclusion Criteria	Pregnancy
Required Medical Information	Diagnosis of one of the following A.) Multiple myeloma that is newly diagnosed, or B.) Erythema nodosum leprosum (ENL)
Age Restrictions	None
Prescriber Restrictions	Prescribed by or in consultation with an oncologist, infectious disease specialist, or dermatologist
Coverage Duration	12 months
Other Criteria	None
Indications	All Medically-accepted Indications.
Off-Label Uses	N/A
Part B Prerequisite	No

Formulary ID: 24449 version 13

Last Updated: 06/20/2024 Effective: 07/01/2024



TIBSOVO

Products Affected

• TIBSOVO

PA Criteria	Criteria Details
Exclusion Criteria	None
Required Medical Information	Diagnosis of one of the following A.) Relapsed or refractory acute myeloid leukemia with a susceptible isocitrate dehydrogenase-1 mutation (as detected by an FDA-approved test), B.) Previously treated, locally advanced or metastatic cholangiocarcinoma with an isocitrate dehydrogenase-1 mutation (as detected by an FDA-approved test.), C.) Acute myeloid leukemia (newly-diagnosed) with susceptible isocitrate dehydrogenase-1 mutation and meets one of the following: 1.) Patient is 75 years of age or older, or 2.) Patient has comorbidities that preclude intensive induction chemotherapy, or D.) Relapsed or refractory myelodysplastic syndromes with a susceptible isocitrate dehydrogenase-1 mutation (as detected by an FDA-approved test)
Age Restrictions	18 years of age and older
Prescriber Restrictions	Prescribed by or in consultation with a gastroenterologist, hematologist, hepatologist, or oncologist
Coverage Duration	12 months
Other Criteria	None
Indications	All Medically-accepted Indications.
Off-Label Uses	N/A
Part B Prerequisite	No

Formulary ID: 24449 version 13 Last Updated: 06/20/2024

Effective: 07/01/2024



TOBI

Products Affected

• TOBI PODHALER

PA Criteria	Criteria Details
Exclusion Criteria	Known sensitivity to any aminoglycoside
Required Medical Information	Diagnosis of cystic fibrosis (confirmed by appropriate diagnostic or genetic testing) and patient has suspected or confirmed Pseudomonas aeruginosa infection in the lungs
Age Restrictions	6 years of age and older
Prescriber Restrictions	None
Coverage Duration	12 months
Other Criteria	None
Indications	All Medically-accepted Indications.
Off-Label Uses	N/A
Part B Prerequisite	No

Formulary ID: 24449 version 13

Last Updated: 06/20/2024 Effective: 07/01/2024



TOLVAPTAN

Products Affected

• tolvaptan

PA Criteria	Criteria Details
Exclusion Criteria	Any of the following A.) Diagnosis of Autosomal Dominant Polycystic Kidney Disease (ADPKD), B.) Urgent need to raise serum sodium acutely, C.) Inability to sense or appropriately respond to thirst, D.) Hypovolemic hyponatremia, E.) Concomitant use of strong CYP 3A Inhibitors (e.g. clarithromycin, ketoconazole, ritonavir), or F.) Anuria
Required Medical Information	Diagnosis of clinically significant hypervolemic or euvolemic hyponatremia (serum sodium less than 125 mEq/L or less marks hyponatremia that is symptomatic and has resisted correction with fluid restriction), including in patients with heart failure and syndrome of inappropriate antidiuretic hormone (SIADH)
Age Restrictions	18 years of age and older
Prescriber Restrictions	None
Coverage Duration	1 month
Other Criteria	None
Indications	All Medically-accepted Indications.
Off-Label Uses	N/A
Part B Prerequisite	No

Formulary ID: 24449 version 13

Last Updated: 06/20/2024 Effective: 07/01/2024



TOPICAL RETINOIDS

Products Affected

• tretinoin external

PA Criteria	Criteria Details
Exclusion Criteria	None
Required Medical Information	Diagnosis of mild to moderate acne vulgaris
Age Restrictions	None
Prescriber Restrictions	None
Coverage Duration	12 months
Other Criteria	None
Indications	All Medically-accepted Indications.
Off-Label Uses	N/A
Part B Prerequisite	No

Formulary ID: 24449 version 13 Last Updated: 06/20/2024

Effective: 07/01/2024



TOREMIFENE

Products Affected

• toremifene citrate

PA Criteria	Criteria Details
Exclusion Criteria	Any of the following A.) Acquired or congenital long QT syndrome, B.) Uncorrected hypokalemia, C.) Uncorrected hypomagnesemia
Required Medical Information	Diagnosis of metastatic breast cancer and patient must have previous inadequate response or intolerance to tamoxifen
Age Restrictions	18 years of age and older
Prescriber Restrictions	Prescribed by or in consultation with an oncologist
Coverage Duration	6 months
Other Criteria	None
Indications	All Medically-accepted Indications.
Off-Label Uses	N/A
Part B Prerequisite	No

Formulary ID: 24449 version 13

Last Updated: 06/20/2024 Effective: 07/01/2024



TRELSTAR

Products Affected

• TRELSTAR MIXJECT

PA Criteria	Criteria Details
Exclusion Criteria	None
Required Medical Information	Diagnosis of advanced prostate cancer
Age Restrictions	None
Prescriber Restrictions	None
Coverage Duration	12 months
Other Criteria	B vs D determination required per CMS guidance
Indications	All Medically-accepted Indications.
Off-Label Uses	N/A
Part B Prerequisite	No

Formulary ID: 24449 version 13 Last Updated: 06/20/2024

Effective: 07/01/2024



TRIENTINE

Products Affected

• trientine hcl

PA Criteria	Criteria Details
Exclusion Criteria	None
Required Medical Information	Diagnosis of Wilson's disease in patients that are intolerant to penicillamine
Age Restrictions	None
Prescriber Restrictions	None
Coverage Duration	12 months
Other Criteria	None
Indications	All Medically-accepted Indications.
Off-Label Uses	N/A
Part B Prerequisite	No

Formulary ID: 24449 version 13

Last Updated: 06/20/2024 Effective: 07/01/2024



TRIKAFTA

Products Affected

• TRIKAFTA

PA Criteria	Criteria Details
Exclusion Criteria	None
Required Medical Information	Diagnosis of cystic fibrosis (CF) and patient has at least 1 F508del mutation in the cystic fibrosis transmembrane conductance regulator (CFTR) gene or a mutation in the CFTR gene that is responsive to elexacaftor/tezacaftor/ivacaftor verified by an FDA-cleared CF mutation test
Age Restrictions	2 years of age and older
Prescriber Restrictions	Prescribed by or in consultation with a pulmonologist or prescribing practitioner is from a CF center accredited by the Cystic Fibrosis Foundation
Coverage Duration	12 months
Other Criteria	None
Indications	All Medically-accepted Indications.
Off-Label Uses	N/A
Part B Prerequisite	No

Formulary ID: 24449 version 13 Last Updated: 06/20/2024

Effective: 07/01/2024



TRUQAP

Products Affected

• TRUQAP

PA Criteria	Criteria Details
Exclusion Criteria	None
Required Medical Information	Diagnosis of hormone receptor (HR)-positive, human epidermal growth factor receptor 2 (HER2)-negative, locally advanced or metastatic breast cancer with 1 or more PIK3CA/AKT1/PTEN-alterations as detected by an FDA-approved test and, A.) patient has had disease progression following 1 or more endocrine-based regimen(s) in the metastatic setting or recurrence on or within 12 months of completing adjuvant therapy, and B.) will be used in combination with fulvestrant injection.
Age Restrictions	18 years of age and older
Prescriber Restrictions	Prescribed by or in consultation with an oncologist or hematologist
Coverage Duration	12 months
Other Criteria	None
Indications	All Medically-accepted Indications.
Off-Label Uses	N/A
Part B Prerequisite	No

Formulary ID: 24449 version 13

Last Updated: 06/20/2024 Effective: 07/01/2024



TUKYSA

Products Affected

• TUKYSA

PA Criteria	Criteria Details
Exclusion Criteria	None
Required Medical Information	Diagnosis of one of the following: A.) advanced unresectable or metastatic HER2-positive breast cancer (including brain metastases) in patients who have received one or more prior anti-HER2-based regimens in the metastatic setting and drug is being used in combination with trastuzumab and capecitabine, or B.) unresectable or metastatic RAS wild-type, HER2-positive colorectal cancer that has progressed following treatment with fluoropyrimidine, oxaliplatin, and irinotecan-based chemotherapy and drug is being used in combination with trastuzumab
Age Restrictions	18 years of age and older
Prescriber Restrictions	Prescribed by or in consultation with an oncologist
Coverage Duration	12 months
Other Criteria	None
Indications	All Medically-accepted Indications.
Off-Label Uses	N/A
Part B Prerequisite	No

Formulary ID: 24449 version 13

Last Updated: 06/20/2024 Effective: 07/01/2024



TURALIO

Products Affected

• TURALIO ORAL CAPSULE 125 MG

PA Criteria	Criteria Details
Exclusion Criteria	None
Required Medical Information	Diagnosis of symptomatic tenosynovial giant cell tumor (TGCT) associated with severe morbidity or functional limitations and not amenable to improvement with surgery
Age Restrictions	18 years of age and older
Prescriber Restrictions	Prescribed by or in consultation with an oncologist
Coverage Duration	12 months
Other Criteria	None
Indications	All Medically-accepted Indications.
Off-Label Uses	N/A
Part B Prerequisite	No

Formulary ID: 24449 version 13

Last Updated: 06/20/2024 Effective: 07/01/2024



TYMLOS

Products Affected

• TYMLOS

PA Criteria	Criteria Details
Exclusion Criteria	None
Required Medical Information	Diagnosis of osteoporosis in men or postmenopausal women and one of the following A.) osteoporotic fracture or multiple risk factors for fracture, or B.) previous trial of/or contraindication to bisphosphonate
Age Restrictions	18 years of age and older
Prescriber Restrictions	None
Coverage Duration	Initial: 12 months, Renewal: 12 months (Maximum 24 month treatment per patient lifetime)
Other Criteria	None
Indications	All Medically-accepted Indications.
Off-Label Uses	N/A
Part B Prerequisite	No

Formulary ID: 24449 version 13

Last Updated: 06/20/2024 Effective: 07/01/2024



UBRELVY

Products Affected

UBRELVY

PA Criteria	Criteria Details
Exclusion Criteria	Concomitant use of strong CYP3A4 inhibitors (e.g., ketoconazole, itraconazole, clarithromycin)
Required Medical Information	Diagnosis of migraine disorder with or without aura and patient has documented trial, inadequate response, or contraindication to at least 1 generic formulary triptan
Age Restrictions	18 years of age and older
Prescriber Restrictions	None
Coverage Duration	12 months
Other Criteria	None
Indications	All Medically-accepted Indications.
Off-Label Uses	N/A
Part B Prerequisite	No

Formulary ID: 24449 version 13

Last Updated: 06/20/2024 Effective: 07/01/2024

 $H9306_24_DRS_09\text{-}001_OE_C$



VALCHLOR

Products Affected

VALCHLOR

PA Criteria	Criteria Details
Exclusion Criteria	None
Required Medical Information	Diagnosis of cutaneous T-cell lymphoma (stage IA and IB mycosis fungoides-type) and patient has received prior skin-directed therapy (e.g. Topical corticosteroids, phototherapy, or topical nitrogen mustard)
Age Restrictions	18 years of age and older
Prescriber Restrictions	Prescribed by or in consultation with an oncologist
Coverage Duration	12 months
Other Criteria	None
Indications	All Medically-accepted Indications.
Off-Label Uses	N/A
Part B Prerequisite	No

Formulary ID: 24449 version 13 Last Updated: 06/20/2024

Effective: 07/01/2024



VANFLYTA

Products Affected

VANFLYTA

PA Criteria	Criteria Details
Exclusion Criteria	None
Required Medical Information	Patient must have all of the following A.) Newly diagnosed acute myeloid leukemia with FLT3-ITD mutation, B.) Used in combination with standard cytarabine and anthracycline induction and cytarabine consolidation, and as maintenance monotherapy following consolidation chemotherapy, and C.) Must be enrolled in the VANFLYTA REMS program
Age Restrictions	18 years of age and older
Prescriber Restrictions	Prescribed by or in consultation with an oncologist or hematologist
Coverage Duration	12 months
Other Criteria	None
Indications	All Medically-accepted Indications.
Off-Label Uses	N/A
Part B Prerequisite	No

Formulary ID: 24449 version 13

Last Updated: 06/20/2024 Effective: 07/01/2024



VENCLEXTA

Products Affected

VENCLEXTA

• VENCLEXTA STARTING PACK

PA Criteria	Criteria Details
Exclusion Criteria	Concomitant use with strong CYP3A inhibitor during the initial and titration phase in patients with CLL or SLL
Required Medical Information	Diagnosis of one of the following A.) chronic lymphocytic leukemia (CLL) or small lymphocytic lymphoma (SLL), or B.) Newly-diagnosed acute myeloid leukemia (AML) and used in combination with azacitidine, decitabine or low-dose cytarabine in patients 75 years or older or who have comorbidities that preclude use of intensive induction chemotherapy
Age Restrictions	18 years of age and older
Prescriber Restrictions	Prescribed by or in consultation with an oncologist or hematologist
Coverage Duration	12 months
Other Criteria	None
Indications	All Medically-accepted Indications.
Off-Label Uses	N/A
Part B Prerequisite	No

Formulary ID: 24449 version 13 Last Updated: 06/20/2024

Effective: 07/01/2024



VERQUVO

Products Affected

VERQUVO

PA Criteria	Criteria Details
Exclusion Criteria	Any of the following A.) Concomitant use of other soluble guanylate cyclase (sGC) stimulators, or B.) Pregnancy
Required Medical Information	Diagnosis of chronic heart failure (HF), NYHA Class II to IV and all of the following 1.) Left ventricular ejection fraction less than 45%, 2.) Previous hospitalization for HF within 6 months or outpatient IV diuretic treatment for HF within 3 months
Age Restrictions	18 years of age and older
Prescriber Restrictions	Prescribed by or in consultation with a cardiologist
Coverage Duration	12 months
Other Criteria	None
Indications	All Medically-accepted Indications.
Off-Label Uses	N/A
Part B Prerequisite	No

Formulary ID: 24449 version 13

Last Updated: 06/20/2024 Effective: 07/01/2024



VERZENIO

Products Affected

VERZENIO

PA Criteria	Criteria Details
Exclusion Criteria	None
Required Medical Information	Diagnosis of one of the following A.) Hormone receptor (HR)-positive, human epidermal growth factor receptor 2 (HER2)-negative, node-positive, early breast cancer and ALL of the following: 1.) Patient is at high risk of recurrence, and 2.) Requested drug will be used in combination with endocrine therapy (tamoxifen or an aromatase inhibitor) for adjuvant treatment, OR B.) Advanced or metastatic, hormone receptor (HR)-positive, human epidermal growth factor receptor 2 (HER2)-negative breast cancer and one of the following 1.) Used in combination with fulvestrant in a patient with disease progression following endocrine therapy, 2.) Used as monotherapy in a patient with disease progression following endocrine therapy and prior chemotherapy in the metastatic setting, or 3.) For postmenopausal women, and men, used as initial endocrine-based treatment in combination with an aromatase inhibitor
Age Restrictions	18 years of age and older
Prescriber Restrictions	Prescribed by or in consultation with an oncologist
Coverage Duration	12 months
Other Criteria	None
Indications	All Medically-accepted Indications.
Off-Label Uses	N/A
Part B Prerequisite	No

Formulary ID: 24449 version 13 Last Updated: 06/20/2024

Effective: 07/01/2024



VIGABATRIN

Products Affected

• vigabatrin

VIGPODER

• VIGADRONE ORAL TABLET

PA Criteria	Criteria Details
Exclusion Criteria	None
Required Medical Information	Diagnosis of one of the following A.) Infantile spasms, or B.) Refractory complex partial seizures and the drug is being used as adjunctive therapy in patients who have responded inadequately to two alternative treatments
Age Restrictions	None
Prescriber Restrictions	Prescribed by or in consultation with a neurologist
Coverage Duration	12 months
Other Criteria	None
Indications	All Medically-accepted Indications.
Off-Label Uses	N/A
Part B Prerequisite	No

Formulary ID: 24449 version 13 Last Updated: 06/20/2024

Effective: 07/01/2024



VIJOICE

Products Affected

• VIJOICE

PA Criteria	Criteria Details
Exclusion Criteria	None
Required Medical Information	Diagnosis of severe manifestations of PIK3CA-Related Overgrowth Spectrum (PROS) in patients who require systemic therapy
Age Restrictions	2 years of age and older
Prescriber Restrictions	None
Coverage Duration	12 months
Other Criteria	None
Indications	All Medically-accepted Indications.
Off-Label Uses	N/A
Part B Prerequisite	No

Formulary ID: 24449 version 13 Last Updated: 06/20/2024

Effective: 07/01/2024



VITRAKVI

Products Affected

VITRAKVI

PA Criteria	Criteria Details
Exclusion Criteria	None
Required Medical Information	Diagnosis of metastatic or surgically unresectable neurotrophic receptor tyrosine kinase (NTRK) gene fusion positive solid tumors without a known acquired resistance mutation and used in patients with unsatisfactory alternative treatments or who have progressed following treatment
Age Restrictions	None
Prescriber Restrictions	Prescribed by or in consultation with an oncologist
Coverage Duration	12 months
Other Criteria	None
Indications	All Medically-accepted Indications.
Off-Label Uses	N/A
Part B Prerequisite	No

Formulary ID: 24449 version 13

Last Updated: 06/20/2024 Effective: 07/01/2024



VIZIMPRO

Products Affected

VIZIMPRO

PA Criteria	Criteria Details
Exclusion Criteria	None
Required Medical Information	Diagnosis of metastatic non-small cell lung cancer with confirmed epidermal growth factor receptor (EGFR) exon 19 deletion or exon 21 L858R substitution mutations as detected by an FDA-approved test
Age Restrictions	18 years of age and older
Prescriber Restrictions	Prescribed by or in consultation with an oncologist
Coverage Duration	12 months
Other Criteria	None
Indications	All Medically-accepted Indications.
Off-Label Uses	N/A
Part B Prerequisite	No

Formulary ID: 24449 version 13 Last Updated: 06/20/2024

Effective: 07/01/2024



VONJO

Products Affected

VONJO

PA Criteria	Criteria Details
Exclusion Criteria	None
Required Medical Information	Diagnosis of intermediate or high-risk primary or secondary myelofibrosis in adults AND a platelet count less than 50 X 10(9)/L
Age Restrictions	None
Prescriber Restrictions	None
Coverage Duration	12 months
Other Criteria	None
Indications	All Medically-accepted Indications.
Off-Label Uses	N/A
Part B Prerequisite	No

Formulary ID: 24449 version 13

Last Updated: 06/20/2024 Effective: 07/01/2024

 $H9306_24_DRS_09\text{-}001_OE_C$



VORICONAZOLE

Products Affected

• voriconazole intravenous

• voriconazole oral

PA Criteria	Criteria Details
Exclusion Criteria	Any of the following A.) Concomitant use of carbamazepine, CYP3A4 substrates (e.g., terfenadine, astemizole, cisapride, pimozide, or quinidine), B.) Concomitant use with high-dose ritonavir (400mg every 12 hours), C.) Concomitant use with ergot alkaloids, D.) Concomitant use with longacting barbiturates, E.) Concomitant use with rifabutin or rifampin, F.) Concomitant use with sirolimus, or G.) Concomitant use with efavirenz at standard doses of 400mg/day or higher
Required Medical Information	Diagnosis of one of the following A.) Invasive aspergillosis, B.) Candidemia, C.) Esophageal Candidiasis, D.) Invasive candidiasis of the skin and abdomen, kidney, bladder wall, and wounds, or E.) Serious fungal infection due to Scedosporium apiospermum or Fusarium species
Age Restrictions	None
Prescriber Restrictions	Prescribed by or in consultation with an infectious disease specialist
Coverage Duration	6 months
Other Criteria	IV formulation: B vs D determination required per CMS guidance
Indications	All Medically-accepted Indications.
Off-Label Uses	N/A
Part B Prerequisite	No

Formulary ID: 24449 version 13 Last Updated: 06/20/2024

Effective: 07/01/2024



VYNDAMAX

Products Affected

VYNDAMAX

PA Criteria	Criteria Details
Exclusion Criteria	None
Required Medical Information	Diagnosis of wild type or hereditary transthyretin related familial amyloid cardiomyopathy
Age Restrictions	18 years of age and older
Prescriber Restrictions	Prescribed by or in consultation with a cardiologist
Coverage Duration	12 months
Other Criteria	None
Indications	All Medically-accepted Indications.
Off-Label Uses	N/A
Part B Prerequisite	No

Formulary ID: 24449 version 13

Last Updated: 06/20/2024 Effective: 07/01/2024



WELIREG

Products Affected

WELIREG

PA Criteria	Criteria Details
Exclusion Criteria	Pregnancy
Required Medical Information	Diagnosis of one of the following A.) Von Hippel-Lindau (VHL) disease and therapy is required for any of the following disease associated tumors that do not require immediate surgery 1.) Renal cell carcinoma (RCC), 2.) Central nervous system (CNS) hemangioblastoma, or 3.) Pancreatic neuroendocrine tumor (pNET), or B.) Advanced renal cell carcinoma following a programmed death receptor-1 or programmed death-ligand 1 inhibitor and a vascular endothelial growth factor tyrosine kinase inhibitor
Age Restrictions	18 years of age and older
Prescriber Restrictions	None
Coverage Duration	12 months
Other Criteria	None
Indications	All Medically-accepted Indications.
Off-Label Uses	N/A
Part B Prerequisite	No

Formulary ID: 24449 version 13

Last Updated: 06/20/2024 Effective: 07/01/2024



XALKORI

Products Affected

XALKORI

PA Criteria	Criteria Details
Exclusion Criteria	None
Required Medical Information	Diagnosis of one of the following A.) Metastatic non-small cell lung cancer (NSCLC) that is anaplastic lymphoma kinase (ALK)-positive or ROS1-positive as detected by an FDA-approved test, B.) Relapsed or refractory systemic anaplastic large cell lymphoma that is anaplastic lymphoma kinase (ALK) positive as detected by an FDA-approved test, or C.) Unresectable, recurrent, or refractory inflammatory myofibroblastic tumors that are anaplastic lymphoma kinase (ALK)-positive
Age Restrictions	None
Prescriber Restrictions	Prescribed by or in consultation with an oncologist
Coverage Duration	12 months
Other Criteria	None
Indications	All Medically-accepted Indications.
Off-Label Uses	N/A
Part B Prerequisite	No

Formulary ID: 24449 version 13

Last Updated: 06/20/2024 Effective: 07/01/2024



XDEMVY

Products Affected

• XDEMVY

PA Criteria	Criteria Details
Exclusion Criteria	None
Required Medical Information	Diagnosis of Demodex blepharitis
Age Restrictions	18 years of age and older
Prescriber Restrictions	None
Coverage Duration	12 months
Other Criteria	None
Indications	All Medically-accepted Indications.
Off-Label Uses	N/A
Part B Prerequisite	No

Formulary ID: 24449 version 13 Last Updated: 06/20/2024

Effective: 07/01/2024



XGEVA

Products Affected

• XGEVA

PA Criteria	Criteria Details
Exclusion Criteria	Hypocalcemia (calcium less than 8.0 mg/dL)
Required Medical Information	Diagnosis of one of the following A.) Bone metastases from a solid tumor and used for the prevention of skeletal related events, B.) Multiple myeloma and used for the prevention of skeletal related events, C.) Hypercalcemia of malignancy refractory to bisphosphonate therapy, or D.) Giant cell tumor of bone that is unresectable or where surgical resection is likely to result in severe morbidity
Age Restrictions	None
Prescriber Restrictions	None
Coverage Duration	12 months
Other Criteria	B vs D determination required per CMS guidance
Indications	All Medically-accepted Indications.
Off-Label Uses	N/A
Part B Prerequisite	No

Formulary ID: 24449 version 13 Last Updated: 06/20/2024

Effective: 07/01/2024



XOLAIR

Products Affected

XOLAIR

PA Criteria	Criteria Details
Exclusion Criteria	None
Required Medical Information	Diagnosis of one of the following A.) Chronic idiopathic urticaria in patients who remain symptomatic despite H1 antihistamine therapy and patient will continue to receive concurrent H1 antihistamine therapy unless contraindicated or not tolerated, B.) Moderate to severe persistent asthma in patients with a positive skin test or in vitro reactivity to a perennial aeroallergen and symptoms are inadequately controlled with inhaled corticosteroids and an additional controller medication (i.e. long acting beta2-agonist, leukotriene modifier, or sustained-release theophylline) and patient has trial and failure, contraindication, or intolerance to Dupixent or Nucala, C.) Nasal polyps in patients with inadequate response to nasal corticosteroids, requested drug will be used as adjunctive treatment, and patient has trial and failure, contraindication, or intolerance to Dupixent, or D.) Reduction of allergic reactions (type I), including anaphylaxis, that may occur with accidental exposure to 1 or more foods in with IgE-mediated food allergy and is being used in conjunction with food allergen avoidance
Age Restrictions	1 year of age and older
Prescriber Restrictions	None
Coverage Duration	12 months
Other Criteria	B vs D determination required per CMS guidance
Indications	All Medically-accepted Indications.
Off-Label Uses	N/A

Formulary ID: 24449 version 13 Last Updated: 06/20/2024

Effective: 07/01/2024



PA Criteria	Criteria Details
Part B Prerequisite	No

Effective: 07/01/2024



XOSPATA

Products Affected

XOSPATA

PA Criteria	Criteria Details
Exclusion Criteria	None
Required Medical Information	Diagnosis of relapsed or refractory acute myeloid leukemia (AML) with a FMS-like tyrosine kinase 3 (FLT3) mutation as detected by an FDA-approved test
Age Restrictions	18 years of age and older
Prescriber Restrictions	Prescribed by or in consultation with an oncologist or hematologist
Coverage Duration	12 months
Other Criteria	None
Indications	All Medically-accepted Indications.
Off-Label Uses	N/A
Part B Prerequisite	No

Formulary ID: 24449 version 13

Last Updated: 06/20/2024 Effective: 07/01/2024



XPOVIO

Products Affected

- XPOVIO (100 MG ONCE WEEKLY) ORAL TABLET THERAPY PACK 50 MG
- XPOVIO (40 MG ONCE WEEKLY) ORAL TABLET THERAPY PACK 40 MG
- XPOVIO (40 MG TWICE WEEKLY)
 ORAL TABLET THERAPY PACK 40
 MG
- XPOVIO (60 MG ONCE WEEKLY) ORAL TABLET THERAPY PACK 60 MG
- XPOVIO (60 MG TWICE WEEKLY)
- XPOVIO (80 MG ONCE WEEKLY)
 ORAL TABLET THERAPY PACK 40
 MG
- XPOVIO (80 MG TWICE WEEKLY)

MG	
PA Criteria	Criteria Details
Exclusion Criteria	None
Required Medical Information	Diagnosis of one of the following A.) Relapsed or refractory multiple myeloma being used in combination with dexamethasone in a patient who has received at least 4 prior therapies and is refractory to at least 2 proteasome inhibitors, at least 2 immunomodulatory agents, and an anti-CD38 monoclonal antibody, B.) Multiple myeloma being used in combination with bortezomib and dexamethasone in a patient who has received at least 1 prior therapy, C.) Relapsed or refractory diffuse large B-cell lymphoma not otherwise specified, or D.) Relapsed or refractory DLBCL arising from follicular lymphoma and patient has received at least 2 lines of systemic therapy
Age Restrictions	18 years of age and older
Prescriber Restrictions	Prescribed by or in consultation with an oncologist or hematologist
Coverage Duration	12 months
Other Criteria	None
Indications	All Medically-accepted Indications.
Off-Label Uses	N/A

Formulary ID: 24449 version 13 Last Updated: 06/20/2024

Effective: 07/01/2024



PA Criteria	Criteria Details
Part B Prerequisite	No

Effective: 07/01/2024



XTANDI

Products Affected

XTANDI

PA Criteria	Criteria Details
Exclusion Criteria	None
Required Medical Information	Diagnosis of one of the following A.) Castration-resistant prostate cancer (CRPC), B.) Metastatic, castration-sensitive prostate cancer (mCSPC). For treatment of CRPC and mCSPC, one of the following applies 1.) Used in combination with a gonadotropin-releasing hormone (GnRH) analog or 2) Patient has received bilateral orchiectomy, or C.) Nonmetastatic castration-sensitive prostate cancer with biochemical recurrence at high risk for metastasis
Age Restrictions	18 years of age and older
Prescriber Restrictions	Prescribed by or in consultation with an oncologist or urologist
Coverage Duration	12 months
Other Criteria	None
Indications	All Medically-accepted Indications.
Off-Label Uses	N/A
Part B Prerequisite	No

Formulary ID: 24449 version 13

Last Updated: 06/20/2024 Effective: 07/01/2024



XURIDEN

Products Affected

• XURIDEN

PA Criteria	Criteria Details
Exclusion Criteria	None
Required Medical Information	Diagnosis of hereditary orotic aciduria
Age Restrictions	None
Prescriber Restrictions	None
Coverage Duration	12 months
Other Criteria	None
Indications	All Medically-accepted Indications.
Off-Label Uses	N/A
Part B Prerequisite	No

Formulary ID: 24449 version 13

Last Updated: 06/20/2024 Effective: 07/01/2024



YONSA

Products Affected

• YONSA

PA Criteria	Criteria Details
Exclusion Criteria	Pregnancy
Required Medical Information	Diagnosis of metastatic, castration-resistant prostate cancer (mCRPC) and used in combination with methylprednisolone. For treatment of mCRPC, one of the following applies: 1.) Used in combination with a gonadotropin-releasing hormone (GnRH) analog or 2) Patient has received bilateral orchiectomy
Age Restrictions	18 years of age and older
Prescriber Restrictions	Prescribed by or in consultation with an oncologist or urologist
Coverage Duration	12 months
Other Criteria	None
Indications	All Medically-accepted Indications.
Off-Label Uses	N/A
Part B Prerequisite	No

Formulary ID: 24449 version 13

Last Updated: 06/20/2024 Effective: 07/01/2024



ZARXIO

Products Affected

• ZARXIO

PA Criteria	Criteria Details
Exclusion Criteria	None
Required Medical Information	Diagnosis of one of the following A.) Chemotherapy induced febrile neutropenia (prophylaxis), B.) Severe chronic neutropenia, C.) Patient is undergoing autologous peripheral-blood progenitor cell transplant to mobilize progenitor cells for collection by leukapheresis, or D.) Hematopoietic subsyndrome of acute radiation syndrome (H-ARS)
Age Restrictions	None
Prescriber Restrictions	None
Coverage Duration	12 months
Other Criteria	None
Indications	All Medically-accepted Indications.
Off-Label Uses	N/A
Part B Prerequisite	No

Formulary ID: 24449 version 13 Last Updated: 06/20/2024

Effective: 07/01/2024



ZEJULA

Products Affected

ZEJULA

PA Criteria	Criteria Details
Exclusion Criteria	None
Required Medical Information	Diagnosis of Advanced or recurrent epithelial ovarian, fallopian tube, or primary peritoneal cancer and used as maintenance therapy in a patient who is in a complete or partial response to platinum-based chemotherapy
Age Restrictions	18 years of age and older
Prescriber Restrictions	Prescribed by or in consultation with an oncologist
Coverage Duration	12 months
Other Criteria	None
Indications	All Medically-accepted Indications.
Off-Label Uses	N/A
Part B Prerequisite	No

Formulary ID: 24449 version 13 Last Updated: 06/20/2024

Effective: 07/01/2024



ZELBORAF

Products Affected

• ZELBORAF

PA Criteria	Criteria Details
Exclusion Criteria	None
Required Medical Information	Diagnosis of one of the following A.) Unresectable or metastatic melanoma and patient has positive BRAF-V600E mutation documented by an FDA-approved test, or B.) Erdheim-Chester disease and patient has documented BRAF V600 mutation
Age Restrictions	18 years of age and older
Prescriber Restrictions	Prescribed by or in consultation with an oncologist or hematologist
Coverage Duration	12 months
Other Criteria	None
Indications	All Medically-accepted Indications.
Off-Label Uses	N/A
Part B Prerequisite	No

Formulary ID: 24449 version 13

Last Updated: 06/20/2024 Effective: 07/01/2024



ZIEXTENZO

Products Affected

ZIEXTENZO

PA Criteria	Criteria Details
Exclusion Criteria	None
Required Medical Information	Diagnosis of a non-myeloid malignancy and drug is being used as prophylaxis for chemotherapy-induced neutropenia
Age Restrictions	None
Prescriber Restrictions	None
Coverage Duration	12 months
Other Criteria	None
Indications	All Medically-accepted Indications.
Off-Label Uses	N/A
Part B Prerequisite	No

Formulary ID: 24449 version 13

Last Updated: 06/20/2024 Effective: 07/01/2024



ZILBRYSQ

Products Affected

ZILBRYSQ

PA Criteria	Criteria Details
Exclusion Criteria	None
Required Medical Information	Diagnosis of generalized myasthenia gravis in adults who are anti- acetylcholine receptor (AChR) antibody positive
Age Restrictions	18 years of age and older
Prescriber Restrictions	None
Coverage Duration	12 months
Other Criteria	None
Indications	All Medically-accepted Indications.
Off-Label Uses	N/A
Part B Prerequisite	No

Formulary ID: 24449 version 13

Last Updated: 06/20/2024 Effective: 07/01/2024



ZOKINVY

Products Affected

ZOKINVY

PA Criteria	Criteria Details
Exclusion Criteria	None
Required Medical Information	Diagnosis of one of the following A.) Hutchinson-Gilford Progeria Syndrome or B.)Processing deficient progeroid laminopathy with documentation of either heterozygous LMNA mutation with progerin-like protein accumulation or homozygous or compound heterozygous ZMPSTE24 mutations
Age Restrictions	None
Prescriber Restrictions	None
Coverage Duration	12 months
Other Criteria	None
Indications	All Medically-accepted Indications.
Off-Label Uses	N/A
Part B Prerequisite	No

Formulary ID: 24449 version 13

Last Updated: 06/20/2024 Effective: 07/01/2024



ZOLINZA

Products Affected

• ZOLINZA

PA Criteria	Criteria Details
Exclusion Criteria	None
Required Medical Information	Diagnosis of primary cutaneous T-cell lymphoma (CTCL) in patients who have progressive, persistent or recurrent disease on or following two systemic therapies (e.g., bexarotene, romidepsin, etc)
Age Restrictions	18 years of age and older
Prescriber Restrictions	None
Coverage Duration	12 months
Other Criteria	None
Indications	All Medically-accepted Indications.
Off-Label Uses	N/A
Part B Prerequisite	No

Formulary ID: 24449 version 13 Last Updated: 06/20/2024

Effective: 07/01/2024



ZTALMY

Products Affected

• ZTALMY

PA Criteria	Criteria Details
Exclusion Criteria	None
Required Medical Information	Diagnosis of seizures associated with cyclin-dependent kinase-like 5 (CDKL5) deficiency disorder (CDD)
Age Restrictions	None
Prescriber Restrictions	Prescribed by or in consultation with a neurologist
Coverage Duration	12 months
Other Criteria	None
Indications	All Medically-accepted Indications.
Off-Label Uses	N/A
Part B Prerequisite	No

Formulary ID: 24449 version 13 Last Updated: 06/20/2024

Effective: 07/01/2024



ZURZUVAE

Products Affected

• ZURZUVAE

PA Criteria	Criteria Details
Exclusion Criteria	None
Required Medical Information	Diagnosis of postpartum depression
Age Restrictions	18 years of age and older
Prescriber Restrictions	None
Coverage Duration	14 days
Other Criteria	None
Indications	All Medically-accepted Indications.
Off-Label Uses	N/A
Part B Prerequisite	No

Formulary ID: 24449 version 13

Last Updated: 06/20/2024 Effective: 07/01/2024



ZYDELIG

Products Affected

• ZYDELIG

PA Criteria	Criteria Details
Exclusion Criteria	History of toxic epidermal necrosis with any drug
Required Medical Information	Diagnosis of Chronic lymphocytic leukemia, used in combination with rituximab and patient has relapsed on at least one prior therapy
Age Restrictions	18 years of age and older
Prescriber Restrictions	Prescribed by or in consultation with an oncologist or hematologist
Coverage Duration	12 months
Other Criteria	None
Indications	All Medically-accepted Indications.
Off-Label Uses	N/A
Part B Prerequisite	No

Formulary ID: 24449 version 13 Last Updated: 06/20/2024

Effective: 07/01/2024



ZYKADIA

Products Affected

• ZYKADIA ORAL TABLET

PA Criteria	Criteria Details
Exclusion Criteria	None
Required Medical Information	Diagnosis of anaplastic lymphoma kinase (ALK)-positive metastatic non-small cell lung cancer (NSCLC)
Age Restrictions	18 years of age and older
Prescriber Restrictions	Prescribed by or in consultation with an oncologist
Coverage Duration	12 months
Other Criteria	None
Indications	All Medically-accepted Indications.
Off-Label Uses	N/A
Part B Prerequisite	No

Formulary ID: 24449 version 13

Last Updated: 06/20/2024 Effective: 07/01/2024



PART B VERSUS PART D

This drug may be covered under Medicare Part B or D depending upon the circumstances. Information may need to be submitted describing the use and setting of the drug to make the determination.

Products Affected

- ABELCET INTRAVENOUS SUSPENSION 5 MG/ML
- acetylcysteine inhalation solution 10 %, 20 %
- acyclovir sodium intravenous solution 50 mg/ml
- albuterol sulfate inhalation nebulization solution (2.5 mg/3ml) 0.083%, 0.63 mg/3ml, 1.25 mg/3ml, 2.5 mg/0.5ml
- amikacin sulfate injection solution 500 mg/2ml
- amphotericin b intravenous solution reconstituted 50 mg
- amphotericin b liposome intravenous suspension reconstituted 50 mg
- ampicillin sodium injection solution reconstituted 1 gm, 125 mg
- ampicillin sodium intravenous solution reconstituted 10 gm
- aprepitant oral capsule 125 mg, 40 mg, 80
 & 125 mg, 80 mg
- azathioprine oral tablet 100 mg, 50 mg, 75 mg
- azithromycin intravenous solution reconstituted 500 mg
- aztreonam injection solution reconstituted
 1 gm, 2 gm
- budesonide inhalation suspension 0.25 mg/2ml, 0.5 mg/2ml, 1 mg/2ml
- calcitonin (salmon) nasal solution 200 unit/act
- calcitriol oral capsule 0.25 mcg, 0.5 mcg
- calcitriol oral solution 1 mcg/ml
- cefotetan disodium injection solution reconstituted 1 gm, 2 gm

Formulary ID: 24449 version 13 Last Updated: 06/20/2024

Effective: 07/01/2024

- cefoxitin sodium intravenous solution reconstituted 1 gm, 10 gm, 2 gm
- cefuroxime sodium injection solution reconstituted 750 mg
- cefuroxime sodium intravenous solution reconstituted 1.5 gm
- chlorpromazine hcl oral concentrate 100 mg/ml, 30 mg/ml
- chlorpromazine hcl oral tablet 10 mg, 100 mg, 200 mg, 25 mg, 50 mg
- cinacalcet hcl oral tablet 30 mg, 60 mg, 90 mg
- ciprofloxacin in d5w intravenous solution 200 mg/100ml
- clindamycin phosphate injection solution 900 mg/6ml
- CLINIMIX E/DEXTROSE (2.75/5) INTRAVENOUS SOLUTION 2.75 %
- CLINIMIX E/DEXTROSE (4.25/10) INTRAVENOUS SOLUTION 4.25 %
- CLINIMIX E/DEXTROSE (4.25/5) INTRAVENOUS SOLUTION 4.25 %
- CLINIMIX E/DEXTROSE (5/15) INTRAVENOUS SOLUTION 5 %
- CLINIMIX E/DEXTROSE (5/20) INTRAVENOUS SOLUTION 5 %
- CLINIMIX/DEXTROSE (4.25/10) INTRAVENOUS SOLUTION 4.25 %
- CLINIMIX/DEXTROSE (4.25/5) INTRAVENOUS SOLUTION 4.25 %
- CLINIMIX/DEXTROSE (5/15) INTRAVENOUS SOLUTION 5 %
- CLINIMIX/DEXTROSE (5/20) INTRAVENOUS SOLUTION 5 %



- colistimethate sodium (cba) injection solution reconstituted 150 mg
- cromolyn sodium inhalation nebulization solution 20 mg/2ml
- cyclophosphamide oral capsule 25 mg, 50 mg
- cyclophosphamide oral tablet 25 mg, 50 mg
- cyclosporine modified oral capsule 100 mg, 25 mg, 50 mg
- cyclosporine modified oral solution 100 mg/ml
- cyclosporine oral capsule 100 mg, 25 mg
- dextrose intravenous solution 10 %, 5 %
- dextrose-sodium chloride intravenous solution 10-0.2 %, 10-0.45 %, 2.5-0.45 %, 5-0.2 %, 5-0.45 %, 5-0.9 %
- diphtheria-tetanus toxoids dt intramuscular suspension 25-5 lfu/0.5ml
- DOXY 100 INTRAVENOUS SOLUTION RECONSTITUTED 100 MG
- ENGERIX-B INJECTION SUSPENSION 20 MCG/ML
- ENGERIX-B INJECTION SUSPENSION PREFILLED SYRINGE 10 MCG/0.5ML, 20 MCG/ML
- ENVARSUS XR ORAL TABLET EXTENDED RELEASE 24 HOUR 0.75 MG, 1 MG, 4 MG
- ERAXIS INTRAVENOUS SOLUTION RECONSTITUTED 100 MG, 50 MG
- ERYTHROCIN LACTOBIONATE INTRAVENOUS SOLUTION RECONSTITUTED 500 MG
- everolimus oral tablet 0.25 mg, 0.5 mg, 0.75 mg, 1 mg
- fluconazole in sodium chloride intravenous solution 200-0.9 mg/100ml-%, 400-0.9 mg/200ml-%
- furosemide injection solution 10 mg/ml
- GENGRAF ORAL CAPSULE 100 MG, 25 MG

H9306_24_DRS_09-001_OE_C

Effective: 07/01/2024

 GENGRAF ORAL SOLUTION 100 MG/ML

- granisetron hcl oral tablet 1 mg
- heparin sodium (porcine) injection solution 1000 unit/ml, 10000 unit/ml, 20000 unit/ml, 5000 unit/ml
- HEPLISAV-B INTRAMUSCULAR SOLUTION PREFILLED SYRINGE 20 MCG/0.5ML
- IMOVAX RABIES INTRAMUSCULAR SUSPENSION RECONSTITUTED 2.5 UNIT/ML
- INTRALIPID INTRAVENOUS EMULSION 20 %, 30 %
- ipratropium bromide inhalation solution 0.02 %
- ipratropium-albuterol inhalation solution 0.5-2.5 (3) mg/3ml
- ISOLYTE-P IN D5W INTRAVENOUS SOLUTION
- ISOLYTE-S PH 7.4 INTRAVENOUS SOLUTION
- kcl in dextrose-nacl intravenous solution 10-5-0.45 meq/l-%-%, 20-5-0.2 meq/l-%-%, 20-5-0.45 meq/l-%-%, 30-5-0.45 meq/l-%-%, 40-5-0.45 meq/l-%-%, 40-5-0.9 meq/l-%-%
- *kcl-lactated ringers-d5w intravenous solution 20 meg/l*
- methotrexate sodium (pf) injection solution 50 mg/2ml
- methotrexate sodium injection solution 50 mg/2ml
- methotrexate sodium oral tablet 2.5 mg
- methylprednisolone oral tablet 16 mg, 32 mg, 4 mg, 8 mg
- metronidazole intravenous solution 500 mg/100ml
- moxifloxacin hcl in nacl intravenous solution 400 mg/250ml
- multiple electro type 1 ph 5.5 intravenous solution



- mycophenolate mofetil oral capsule 250 mg
- mycophenolate mofetil oral suspension reconstituted 200 mg/ml
- mycophenolate mofetil oral tablet 500 mg
- mycophenolate sodium oral tablet delayed release 180 mg, 360 mg
- nafcillin sodium injection solution reconstituted 1 gm, 2 gm
- nafcillin sodium intravenous solution reconstituted 10 gm
- NUTRILIPID INTRAVENOUS EMULSION 20 %
- ondansetron hcl oral solution 4 mg/5ml
- ondansetron hcl oral tablet 4 mg, 8 mg
- ondansetron oral tablet dispersible 4 mg, 8 mg
- oxacillin sodium in dextrose intravenous solution 1 gm/50ml, 2 gm/50ml
- oxacillin sodium injection solution reconstituted 1 gm, 2 gm
- oxacillin sodium intravenous solution reconstituted 10 gm
- PANZYGA INTRAVENOUS SOLUTION 1 GM/10ML, 10 GM/100ML, 2.5 GM/25ML, 20 GM/200ML, 30 GM/300ML, 5 GM/50ML
- paricalcitol oral capsule 1 mcg, 2 mcg, 4 mcg
- penicillin g potassium injection solution reconstituted 20000000 unit
- penicillin g sodium injection solution reconstituted 5000000 unit
- pentamidine isethionate inhalation solution reconstituted 300 mg
- pentamidine isethionate injection solution reconstituted 300 mg
- perphenazine oral tablet 4 mg, 8 mg
- PLASMA-LYTE A INTRAVENOUS SOLUTION

- potassium chloride in nacl intravenous solution 20-0.45 meq/l-%, 20-0.9 meq/l-%, 40-0.9 meg/l-%
- potassium chloride intravenous solution 10 meq/100ml, 2 meq/ml, 2 meq/ml (20 ml), 20 meq/100ml, 40 meq/100ml
- potassium cl in dextrose 5% intravenous solution 20 meg/l
- prednisolone oral solution 15 mg/5ml
- prednisolone sodium phosphate oral solution 10 mg/5ml, 20 mg/5ml, 25 mg/5ml, 6.7 (5 base) mg/5ml
- prednisolone sodium phosphate oral tablet dispersible 10 mg, 15 mg, 30 mg
- PREDNISONE INTENSOL ORAL CONCENTRATE 5 MG/ML
- prednisone oral solution 5 mg/5ml
- prednisone oral tablet 1 mg, 10 mg, 2.5 mg, 20 mg, 5 mg, 50 mg
- PREHEVBRIO INTRAMUSCULAR SUSPENSION 10 MCG/ML
- PREMASOL INTRAVENOUS SOLUTION 10 %
- PRIVIGEN INTRAVENOUS SOLUTION 20 GM/200ML
- prochlorperazine maleate oral tablet 10 mg, 5 mg
- PROGRAF ORAL PACKET 0.2 MG, 1 MG
- PROSOL INTRAVENOUS SOLUTION 20 %
- PULMOZYME INHALATION SOLUTION 2.5 MG/2.5ML
- RABAVERT INTRAMUSCULAR SUSPENSION RECONSTITUTED
- RECOMBIVAX HB INJECTION SUSPENSION 10 MCG/ML, 40 MCG/ML, 5 MCG/0.5ML
- RECOMBIVAX HB INJECTION SUSPENSION PREFILLED SYRINGE 10 MCG/ML, 5 MCG/0.5ML
- sirolimus oral solution 1 mg/ml

Effective: 07/01/2024



- sirolimus oral tablet 0.5 mg, 1 mg, 2 mg
- tacrolimus oral capsule 0.5 mg, 1 mg, 5 mg
- TDVAX INTRAMUSCULAR SUSPENSION 2-2 LF/0.5ML
- TEFLARO INTRAVENOUS SOLUTION RECONSTITUTED 400 MG, 600 MG
- TENIVAC INTRAMUSCULAR INJECTABLE 5-2 LFU, 5-2 LFU (INJECTION)
- tigecycline intravenous solution reconstituted 50 mg
- tobramycin inhalation nebulization solution 300 mg/5ml

- tobramycin sulfate injection solution 10 mg/ml, 80 mg/2ml
- TPN ELECTROLYTES INTRAVENOUS CONCENTRATE
- TRAVASOL INTRAVENOUS SOLUTION 10 %
- TREXALL ORAL TABLET 10 MG, 15 MG, 5 MG, 7.5 MG
- TROPHAMINE INTRAVENOUS SOLUTION 10 %
- VARUBI (180 MG DOSE) ORAL TABLET THERAPY PACK 2 X 90 MG
- XATMEP ORAL SOLUTION 2.5 MG/ML

Effective: 07/01/2024



Index

\mathbf{A}	AVONEX PEN INTRAMUSCULAR
ABELCET INTRAVENOUS	AUTO-INJECTOR KIT 144
SUSPENSION 5 MG/ML271	AVONEX PREFILLED
abiraterone acetate 1	INTRAMUSCULAR PREFILLED
acetylcysteine inhalation solution 10 %, 20	SYRINGE KIT 144
% 271	AYVAKIT 16
acitretin2	azathioprine oral tablet 100 mg, 50 mg, 75
ACTIMMUNE3	mg 271
acyclovir sodium intravenous solution 50	azithromycin intravenous solution
mg/ml 271	reconstituted 500 mg271
ADEMPAS 4	aztreonam injection solution reconstituted 1
AKEEGA 5	gm, 2 gm 271
albuterol sulfate inhalation nebulization	В
solution (2.5 mg/3ml) 0.083%, 0.63	BALVERSA17
mg/3ml, 1.25 mg/3ml, 2.5 mg/0.5ml 271	BENLYSTA SUBCUTANEOUS 18
ALECENSA6	BESREMI 19
ALUNBRIG8	BETASERON SUBCUTANEOUS KIT 144
ALVAIZ9	bexarotene external20
ambrisentan10	bexarotene oral21
amikacin sulfate injection solution 500	bosentan22
mg/2ml271	BOSULIF23
amphotericin b intravenous solution	BRAFTOVI ORAL CAPSULE 75 MG 24
reconstituted 50 mg271	BRONCHITOL25
amphotericin b liposome intravenous	BRUKINSA26
suspension reconstituted 50 mg 271	budesonide inhalation suspension 0.25
ampicillin sodium injection solution	mg/2ml, 0.5 mg/2ml, 1 mg/2ml 271
reconstituted 1 gm, 125 mg271	BYLVAY27
ampicillin sodium intravenous solution	BYLVAY (PELLETS)27
reconstituted 10 gm271	C
aprepitant oral capsule 125 mg, 40 mg, 80 &	CABLIVI
125 mg, 80 mg 271	CABOMETYX29
ARCALYST11	calcitonin (salmon) nasal solution 200
ARIKAYCE12	unit/act271
armodafinil35	calcitriol oral capsule 0.25 mcg, 0.5 mcg 271
AUGTYRO13	calcitriol oral solution 1 mcg/ml271
AURYXIA14	CALQUENCE30
AUSTEDO15	CAMZYOS31
AUSTEDO XR 15	CAPRELSA32
AUSTEDO XR PATIENT TITRATION . 15	carglumic acid oral tablet soluble 33
Formulary ID: 24449 version 13	
Last Updated: 06/20/2024	
Effective: 07/01/2024	
H9306_24_DRS_09-001_OE_C	
11/300_4+_DNS_0/-001_OD_C	



CAYSTON34	COMETRIQ (140 MG DAILY DOSE)
cefotetan disodium injection solution	ORAL KIT 3 X 20 MG & 80 MG 36
reconstituted 1 gm, 2 gm271	COMETRIQ (60 MG DAILY DOSE) 36
cefoxitin sodium intravenous solution	COPAXONE SUBCUTANEOUS
reconstituted 1 gm, 10 gm, 2 gm 271	SOLUTION PREFILLED SYRINGE 84
cefuroxime sodium injection solution	COPIKTRA37
reconstituted 750 mg271	CORLANOR ORAL TABLET 38
cefuroxime sodium intravenous solution	COSENTYX (300 MG DOSE) 39, 40
reconstituted 1.5 gm271	COSENTYX SENSOREADY (300 MG) 39,
chlorpromazine hcl oral concentrate 100	40
mg/ml, 30 mg/ml271	COSENTYX SUBCUTANEOUS
chlorpromazine hcl oral tablet 10 mg, 100	SOLUTION PREFILLED SYRINGE 75
mg, 200 mg, 25 mg, 50 mg271	MG/0.5ML39, 40
cinacalcet hcl oral tablet 30 mg, 60 mg, 90	COSENTYX UNOREADY39, 40
mg271	COTELLIC41
ciprofloxacin in d5w intravenous solution	cromolyn sodium inhalation nebulization
200 mg/100ml 271	solution 20 mg/2ml 272
clindamycin phosphate injection solution	cyclophosphamide oral capsule 25 mg, 50
900 mg/6ml271	mg 272
CLINIMIX E/DEXTROSE (2.75/5)	cyclophosphamide oral tablet 25 mg, 50 mg
INTRAVENOUS SOLUTION 2.75 % 271	272
CLINIMIX E/DEXTROSE (4.25/10)	cyclosporine modified oral capsule 100 mg,
INTRAVENOUS SOLUTION 4.25 % 271	25 mg, 50 mg 272
CLINIMIX E/DEXTROSE (4.25/5)	cyclosporine modified oral solution 100
INTRAVENOUS SOLUTION 4.25 % 271	mg/ml
CLINIMIX E/DEXTROSE (5/15)	cyclosporine oral capsule 100 mg, 25 mg272
INTRAVENOUS SOLUTION 5 % 271	CYSTADROPS43
CLINIMIX E/DEXTROSE (5/20)	CYSTAGON42
INTRAVENOUS SOLUTION 5 % 271	CYSTARAN
CLINIMIX/DEXTROSE (4.25/10)	D
INTRAVENOUS SOLUTION 4.25 % 271	dalfampridine er
CLINIMIX/DEXTROSE (4.25/5)	DAURISMO45
INTRAVENOUS SOLUTION 4.25 % 271	DAYBUE46
CLINIMIX/DEXTROSE (5/15)	deferasirox granules47
INTRAVENOUS SOLUTION 5 % 271	deferasirox oral tablet47
CLINIMIX/DEXTROSE (5/20)	deferasirox oral tablet soluble47
INTRAVENOUS SOLUTION 5 % 271	deferiprone
colistimethate sodium (cba) injection	dextrose intravenous solution 10 %, 5 % 272
solution reconstituted 150 mg 272	dextrose-sodium chloride intravenous
COMETRIQ (100 MG DAILY DOSE)	solution 10-0.2 %, 10-0.45 %, 2.5-0.45 %,
ORAL KIT 80 & 20 MG 36	5-0.2 %, 5-0.45 %, 5-0.9 %
	DIACOMIT49

Effective: 07/01/2024



diclofenac sodium external gel 3 % 50	everolimus oral tablet 0.25 mg, 0.5 mg, 0.75
DIFICID51	mg, 1 mg272
dimethyl fumarate oral 52	everolimus oral tablet 10 mg, 2.5 mg, 5 mg,
dimethyl fumarate starter pack oral capsule	7.5 mg
delayed release therapy pack 52	everolimus oral tablet soluble
diphtheria-tetanus toxoids dt intramuscular	EVRYSDI 68
suspension 25-5 lfu/0.5ml	EXKIVITY69
DOJOLVI 53	F
DOXY 100 INTRAVENOUS SOLUTION	febuxostat70
RECONSTITUTED 100 MG 272	fentanyl72
dronabinol54	fentanyl citrate buccal lozenge on a handle
droxidopa 55	71
DUPIXENT56	FERRIPROX ORAL SOLUTION 48
\mathbf{E}	FERRIPROX TWICE-A-DAY48
ELIGARD126	FILSPARI73
EMGALITY 57	fingolimod hcl74
ENBREL MINI 58	FINTEPLA75
ENBREL SUBCUTANEOUS SOLUTION	FIRMAGON (240 MG DOSE)76
25 MG/0.5ML 58	FIRMAGON SUBCUTANEOUS
ENBREL SUBCUTANEOUS SOLUTION	SOLUTION RECONSTITUTED 80 MG
PREFILLED SYRINGE58	76
ENBREL SURECLICK SUBCUTANEOUS	fluconazole in sodium chloride intravenous
SOLUTION AUTO-INJECTOR 58	solution 200-0.9 mg/100ml-%, 400-0.9
ENDARI 59	mg/200ml-%272
ENGERIX-B INJECTION SUSPENSION	FOTIVDA77
20 MCG/ML 272	FRUZAQLA78
ENGERIX-B INJECTION SUSPENSION	furosemide injection solution 10 mg/ml. 272
PREFILLED SYRINGE 10 MCG/0.5ML,	G
20 MCG/ML272	GALAFOLD79
ENSPRYNG 60	GATTEX 80
ENVARSUS XR ORAL TABLET	GAVRETO 81
EXTENDED RELEASE 24 HOUR 0.75	gefitinib82
MG, 1 MG, 4 MG272	GENGRAF ORAL CAPSULE 100 MG, 25
EPIDIOLEX61	MG272
ERAXIS INTRAVENOUS SOLUTION	GENGRAF ORAL SOLUTION 100
RECONSTITUTED 100 MG, 50 MG 272	MG/ML272
ERIVEDGE	GILOTRIF 83
ERLEADA	GLEOSTINE ORAL CAPSULE 10 MG,
erlotinib hcl65	100 MG, 40 MG 85
ERYTHROCIN LACTOBIONATE	granisetron hel oral tablet 1 mg 272
INTRAVENOUS SOLUTION	
RECONSTITUTED 500 MG272	
Formulary ID: 24449 version 13	
Last Updated: 06/20/2024	
Effective: 07/01/2024	
H9306_24_DRS_09-001_OE_C	



H	INCRELEX 100
heparin sodium (porcine) injection solution	INLYTA101
1000 unit/ml, 10000 unit/ml, 20000	INQOVI 102
unit/ml, 5000 unit/ml 272	INREBIC
HEPLISAV-B INTRAMUSCULAR	INTRALIPID INTRAVENOUS
SOLUTION PREFILLED SYRINGE 20	EMULSION 20 %, 30 % 272
MCG/0.5ML	INTRAROSA104
HUMIRA (2 PEN)	ipratropium bromide inhalation solution
HUMIRA (2 SYRINGE)	0.02 %
SUBCUTANEOUS PREFILLED	ipratropium-albuterol inhalation solution
SYRINGE KIT 10 MG/0.1ML, 20	0.5-2.5 (3) mg/3ml
MG/0.2ML, 40 MG/0.4ML, 40	ISOLYTE-P IN D5W INTRAVENOUS
MG/0.8ML 90, 91	SOLUTION
HUMIRA-CD/UC/HS STARTER	ISOLYTE-S PH 7.4 INTRAVENOUS
SUBCUTANEOUS PEN-INJECTOR	SOLUTION272
KIT 80 MG/0.8ML90, 91	ISTURISA ORAL TABLET 1 MG, 5 MG
HUMIRA-PED<40KG CROHNS	
STARTER90, 91	itraconazole oral
HUMIRA-PED>/=40KG CROHNS START	ivermectin oral
90, 91	IWILFIN
HUMIRA-PED>/=40KG UC STARTER 90,	J
91	JAKAFI110
HUMIRA-PS/UV/ADOL HS STARTER 90,	JAYPIRCA111
91	JOENJA
HUMIRA-PSORIASIS/UVEIT STARTER	JUXTAPID ORAL CAPSULE 10 MG, 20
90, 91	MG, 30 MG, 5 MG 113
HYFTOR	K
T	KALYDECO114
IBRANCE	kcl in dextrose-nacl intravenous solution 10-
icatibant acetate subcutaneous solution	5-0.45 meq/l-%-%, 20-5-0.2 meq/l-%-%,
prefilled syringe	20-5-0.45 meg/l-%-%, 20-5-0.9 meg/l-%-
ICLUSIG	%, 30-5-0.45 meq/1-%-%, 40-5-0.45
IDHIFA96	meq/1-%-%, 40-5-0.9 meq/1-%-% 272
imatinib mesylate	kcl-lactated ringers-d5w intravenous
IMBRUVICA ORAL CAPSULE 98	solution 20 meq/l272
IMBRUVICA ORAL SUSPENSION 98	KESIMPTA
IMBRUVICA ORAL TABLET 140 MG,	KINERET SUBCUTANEOUS SOLUTION
280 MG, 420 MG	PREFILLED SYRINGE
IMOVAX RABIES INTRAMUSCULAR	KISQALI (200 MG DOSE) 117, 118
SUSPENSION RECONSTITUTED 2.5	KISQALI (400 MG DOSE) 117, 118
UNIT/ML	KISQALI (600 MG DOSE)
INBRIJA	KISQALI FEMARA (200 MG DOSE) 119
Formulary ID: 24449 version 13	
Last Updated: 06/20/2024	
Effective: 07/01/2024	
H9306_24_DRS_09-001_OE_C	



KISQALI FEMARA (400 MG DOSE) 119	M
KISQALI FEMARA (600 MG DOSE) 119	MATULANE 138
KOSELUGO 120	MAVYRET 89
KRAZATI121	MAYZENT139
\mathbf{L}	MAYZENT STARTER PACK 139
lapatinib ditosylate 122	MEKINIST 140
lenalidomide123	MEKTOVI 141
LENVIMA (10 MG DAILY DOSE) 124	methotrexate sodium (pf) injection solution
LENVIMA (12 MG DAILY DOSE) 124	50 mg/2ml
LENVIMA (14 MG DAILY DOSE) 124	methotrexate sodium injection solution 50
LENVIMA (18 MG DAILY DOSE) 124	mg/2ml 272
LENVIMA (20 MG DAILY DOSE) 124	methotrexate sodium oral tablet 2.5 mg 272
LENVIMA (24 MG DAILY DOSE) 124	methylprednisolone oral tablet 16 mg, 32
LENVIMA (4 MG DAILY DOSE) 124	mg, 4 mg, 8 mg272
LENVIMA (8 MG DAILY DOSE) 124	metronidazole intravenous solution 500
LEUKINE INJECTION SOLUTION	mg/100ml272
RECONSTITUTED125	mifepristone oral tablet 300 mg 142
leuprolide acetate (3 month) 126	miglustat143
leuprolide acetate injection 126	modafinil oral35
lidocaine external patch 5 % 127	MOUNJARO 86
linezolid intravenous solution 600 mg/300ml	moxifloxacin hel in nacl intravenous
128	solution 400 mg/250ml 272
linezolid oral tablet	multiple electro type 1 ph 5.5 intravenous
LIVMARLI	solution272
LIVTENCITY	mycophenolate mofetil oral capsule 250 mg
LONSURF	273
LORBRENA 132	mycophenolate mofetil oral suspension
LUMAKRAS 133	reconstituted 200 mg/ml 273
LUPKYNIS	mycophenolate mofetil oral tablet 500 mg
LUPRON DEPOT (1-MONTH)	273
LUPRON DEPOT (3-MONTH)	mycophenolate sodium oral tablet delayed
LUPRON DEPOT (4-MONTH)	release 180 mg, 360 mg 273
LUPRON DEPOT (6-MONTH)	N
LUPRON DEPOT-PED (1-MONTH)	nafcillin sodium injection solution
INTRAMUSCULAR KIT 7.5 MG 126	reconstituted 1 gm, 2 gm
LUPRON DEPOT-PED (3-MONTH)	nafcillin sodium intravenous solution
INTRAMUSCULAR KIT 11.25 MG. 126	reconstituted 10 gm
LUPRON DEPOT-PED (6-MONTH) 126	NAMZARIC
LYNPARZA ORAL TABLET 135, 136	NERLYNX 146
LYTGOBI (12 MG DAILY DOSE) 137	NINLARO
LYTGOBI (16 MG DAILY DOSE) 137	nitisinone
LYTGOBI (20 MG DAILY DOSE) 137	NOXAFIL ORAL PACKET 173
Formulary ID: 24449 version 13 Last Updated: 06/20/2024 Effective: 07/01/2024 H9306_24_DRS_09-001_OE_C	



NUBEQA149	OZEMPIC (2 MG/DOSE) 86
NUCALA	P
NUEDEXTA151	PANRETIN 165
NUPLAZID ORAL CAPSULE 152	PANZYGA INTRAVENOUS SOLUTION
NUPLAZID ORAL TABLET 10 MG 152	1 GM/10ML, 10 GM/100ML, 2.5
NUTRILIPID INTRAVENOUS	GM/25ML, 20 GM/200ML, 30
EMULSION 20 %	GM/300ML, 5 GM/50ML 273
0	paricalcitol oral capsule 1 mcg, 2 mcg, 4
octreotide acetate injection solution 100	mcg273
mcg/ml, 1000 mcg/ml, 200 mcg/ml, 50	pazopanib hcl 166
mcg/ml, 500 mcg/ml 153	PEGASYS SUBCUTANEOUS SOLUTION
ODOMZO154	180 MCG/ML 167
OFEV 155	PEGASYS SUBCUTANEOUS SOLUTION
OGSIVEO ORAL TABLET 50 MG 156	PREFILLED SYRINGE167
OJJAARA 157	PEMAZYRE168
OMNITROPE SUBCUTANEOUS	penicillin g potassium injection solution
SOLUTION CARTRIDGE 87, 88	reconstituted 20000000 unit
OMNITROPE SUBCUTANEOUS	penicillin g sodium injection solution
SOLUTION RECONSTITUTED 87, 88	reconstituted 5000000 unit
ondansetron hcl oral solution 4 mg/5ml 273	pentamidine isethionate inhalation solution
ondansetron hcl oral tablet 4 mg, 8 mg 273	reconstituted 300 mg273
ondansetron oral tablet dispersible 4 mg, 8	pentamidine isethionate injection solution
mg273	reconstituted 300 mg273
ONUREG	perphenazine oral tablet 4 mg, 8 mg 273
OPSUMIT	PIQRAY (200 MG DAILY DOSE) 169
ORGOVYX160	PIQRAY (250 MG DAILY DOSE) 169
ORKAMBI	PIQRAY (300 MG DAILY DOSE) 169
ORSERDU	pirfenidone
OSPHENA	PLASMA-LYTE A INTRAVENOUS
OTEZLA	SOLUTION
oxacillin sodium in dextrose intravenous	POMALYST171
solution 1 gm/50ml, 2 gm/50ml 273	posaconazole oral
oxacillin sodium injection solution	potassium chloride in nacl intravenous
reconstituted 1 gm, 2 gm273	solution 20-0.45 meq/l-%, 20-0.9 meq/l-
oxacillin sodium intravenous solution	%, 40-0.9 meq/l-%273
reconstituted 10 gm	potassium chloride intravenous solution 10
OZEMPIC (0.25 OR 0.5 MG/DOSE)	meq/100ml, 2 meq/ml, 2 meq/ml (20 ml),
SUBCUTANEOUS SOLUTION PEN-	20 meq/100ml, 40 meq/100ml
INJECTOR 2 MG/3ML 86	potassium cl in dextrose 5% intravenous
OZEMPIC (1 MG/DOSE)	solution 20 meq/l273
SUBCUTANEOUS SOLUTION PEN-	prednisolone oral solution 15 mg/5ml 273
INJECTOR 4 MG/3ML 86	predimental order solution 15 mg/smi 275
Formulary ID: 24449 version 13	
Last Updated: 06/20/2024	
Effective: 07/01/2024	
H9306_24_DRS_09-001_OE_C	



prednisolone sodium phosphate oral solution	REPATHA	180
10 mg/5ml, 20 mg/5ml, 25 mg/5ml, 6.7 (5	REPATHA PUSHTRONEX SYSTEM	180
base) mg/5ml273	REPATHA SURECLICK	180
prednisolone sodium phosphate oral tablet	RETACRIT INJECTION SOLUTION	
dispersible 10 mg, 15 mg, 30 mg 273	10000 UNIT/ML, 10000	
PREDNISONE INTENSOL ORAL	UNIT/ML(1ML), 2000 UNIT/ML, 20	000
CONCENTRATE 5 MG/ML273	UNIT/ML, 3000 UNIT/ML, 4000	
prednisone oral solution 5 mg/5ml 273	UNIT/ML, 40000 UNIT/ML	62
prednisone oral tablet 1 mg, 10 mg, 2.5 mg,	RETEVMO	
20 mg, 5 mg, 50 mg273	REZLIDHIA	
PREHEVBRIO INTRAMUSCULAR	REZUROCK	
SUSPENSION 10 MCG/ML 273	riluzole	
PREMASOL INTRAVENOUS SOLUTION	RINVOQ	
10 % 273	RIVFLOZA	
PREVYMIS ORAL 174	ROZLYTREK	
PRIVIGEN INTRAVENOUS SOLUTION	RUBRACA	
20 GM/200ML273	RYBELSUS	
prochlorperazine maleate oral tablet 10 mg,	RYDAPT	
5 mg	S	
PROGRAF ORAL PACKET 0.2 MG, 1 MG	sapropterin dihydrochloride oral packet.	190
273	sapropterin dihydrochloride oral tablet	
PROLASTIN-C INTRAVENOUS	SCEMBLIX	
SOLUTION7	SIGNIFOR	
PROMACTA 175	sildenafil citrate oral tablet 20 mg	193
PROSOL INTRAVENOUS SOLUTION 20	sirolimus oral solution 1 mg/ml	
% 273	sirolimus oral tablet 0.5 mg, 1 mg, 2 mg	
PULMOZYME INHALATION	SIRTURO	
SOLUTION 2.5 MG/2.5ML273	SKYRIZI PEN	195
Q	SKYRIZI SUBCUTANEOUS	195
QINLOCK 176	sodium oxybate	196
quinine sulfate oral 177	sofosbuvir-velpatasvir	89
R	SOHONOS	
RABAVERT INTRAMUSCULAR	SOLTAMOX	198
SUSPENSION RECONSTITUTED 273	SOMAVERT	199
RAVICTI 178	sorafenib tosylate	200
RECOMBIVAX HB INJECTION	SPRYCEL	201
SUSPENSION 10 MCG/ML, 40	STELARA SUBCUTANEOUS	
MCG/ML, 5 MCG/0.5ML 273	SOLUTION 45 MG/0.5ML	202
RECOMBIVAX HB INJECTION	STELARA SUBCUTANEOUS	
SUSPENSION PREFILLED SYRINGE	SOLUTION PREFILLED SYRINGE	202
10 MCG/ML, 5 MCG/0.5ML 273	STIVARGA	203
REGRANEX179	sunitinib malate	204
Formulary ID: 24449 version 13 Last Updated: 06/20/2024 Effective: 07/01/2024 H9306_24_DRS_09-001_OE_C		



SYMDEKO205	tobramycin sulfate injection solution 10
SYMLINPEN 120 SUBCUTANEOUS	mg/ml, 80 mg/2ml274
SOLUTION PEN-INJECTOR206	tolvaptan225
SYMLINPEN 60 SUBCUTANEOUS	toremifene citrate
SOLUTION PEN-INJECTOR 206	TPN ELECTROLYTES INTRAVENOUS
SYNAREL	CONCENTRATE274
T	TRAVASOL INTRAVENOUS SOLUTION
TABLOID	10 %
TABRECTA	TRELSTAR MIXJECT228
tacrolimus oral capsule 0.5 mg, 1 mg, 5 mg	tretinoin external
274	TREXALL ORAL TABLET 10 MG, 15
TAFINLAR	MG, 5 MG, 7.5 MG 274
TAGRISSO211	trientine hcl
TAKHZYRO212	TRIKAFTA230
TALZENNA	TROPHAMINE INTRAVENOUS
TASIGNA214	SOLUTION 10 %
TAVNEOS	TRULICITY86
tazarotene external cream	TRUQAP231
tazarotene external gel	TUKYSA
TAZORAC EXTERNAL CREAM 0.05 %	TURALIO ORAL CAPSULE 125 MG 233
216	TYMLOS
TAZVERIK217	U
TDVAX INTRAMUSCULAR	UBRELVY235
SUSPENSION 2-2 LF/0.5ML 274	V
TEFLARO INTRAVENOUS SOLUTION	VALCHLOR236
RECONSTITUTED 400 MG, 600 MG	VANFLYTA
274	VARUBI (180 MG DOSE) ORAL
TEGSEDI	TABLET THERAPY PACK 2 X 90 MG
TENIVAC INTRAMUSCULAR	
INJECTABLE 5-2 LFU, 5-2 LFU	VENCLEXTA238
(INJECTION)274	VENCLEXTA STARTING PACK 238
TEPMETKO	VERQUVO239
teriparatide (recombinant) subcutaneous	VERZENIO239
solution pen-injector 620 mcg/2.48ml 220	VERZENIO240 VICTOZA SUBCUTANEOUS SOLUTION
tetrabenazine	PEN-INJECTOR86
THALOMID	
TIBSOVO	vigabatrin 241 VIGADRONE ORAL TABLET 241
tigecycline intravenous solution	VIGPODER
reconstituted 50 mg	VIJOICE
TOBI PODHALER	VITRAKVI
tobramycin inhalation nebulization solution	VIZIMPRO
300 mg/5ml 274	VONJO245
Formulary ID: 24449 version 13	
Last Updated: 06/20/2024	
Effective: 07/01/2024	



voriconazole intravenous 246	XPOVIO (60 MG TWICE WEEKLY) 255,
voriconazole oral246	256
VOSEVI89	XPOVIO (80 MG ONCE WEEKLY) ORAL
VYNDAMAX247	TABLET THERAPY PACK 40 MG. 255,
\mathbf{W}	256
WELIREG248	XPOVIO (80 MG TWICE WEEKLY) 255,
X	256
XALKORI249	XTANDI257
XATMEP ORAL SOLUTION 2.5 MG/ML	XURIDEN258
274	Y
XDEMVY250	YARGESA143
XGEVA251	YONSA259
XOLAIR252, 253	${f Z}$
XOSPATA	ZARXIO260
XPOVIO (100 MG ONCE WEEKLY)	ZEJULA261
ORAL TABLET THERAPY PACK 50	ZELBORAF262
MG255, 256	ZIEXTENZO263
XPOVIO (40 MG ONCE WEEKLY) ORAL	ZILBRYSQ264
TABLET THERAPY PACK 40 MG. 255,	ZOKINVY265
256	ZOLINZA266
XPOVIO (40 MG TWICE WEEKLY)	ZTALMY267
ORAL TABLET THERAPY PACK 40	ZURZUVAE268
MG255, 256	ZYDELIG269
XPOVIO (60 MG ONCE WEEKLY) ORAL	ZYKADIA ORAL TABLET 270
TABLET THERAPY PACK 60 MG. 255,	
256	

Effective: 07/01/2024